

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 January 2003 (16.01.2003)

PCT

(10) International Publication Number
WO 03/004630 A2

(51) International Patent Classification⁷: C12N 9/00

(21) International Application Number: PCT/EP02/07520

(22) International Filing Date: 5 July 2002 (05.07.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
01116412.6 6 July 2001 (06.07.2001) EP
60/379,412 13 May 2002 (13.05.2002) US

(71) Applicant (*for all designated States except US*):
**ARBEITSGEMEINSCHAFT DEUTSCHER
RINDERZÜCHTER E.V. (ADR)** [DE/DE]; Ade-
nauerallee 174, 53113 Bonn (DE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **FRIES,**
Hans-Rudolf [CH/DE]; Philipp-Dirr-Strasse 28, 85354
Freising (DE). **WINTER, Andreas** [DE/DE]; Brunnhaus-
gasse 3, 85354 Freising (DE).

(74) Agent: **VOSSIUS & PARTNER**; Siebertstrasse 4, 81675
Munich (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: METHOD OF TESTING A MAMMAL FOR ITS PREDISPOSITION FOR FAT CONTENT OF MILK AND/OR ITS
PREDISPOSITION FOR MEAT MARBLING

(57) Abstract: The present invention relates to a newly identified nucleic acid sequence of an allele of the polymorphic bovine
DGAT gene. Moreover, the present invention relates to a method of testing a mammal for its predisposition for fat content of milk
and/or its predisposition for meat marbling.

WO 03/004630 A2

THIS PAGE BLANK (USPTO)

Method of testing a mammal for its predisposition for fat content of milk and/or its predisposition for meat marbling

The present invention relates to a newly identified nucleic acid sequence of an allele of the polymorphic bovine *DGAT* gene. Moreover, the present invention relates to a method of testing a mammal for its predisposition for fat content of milk and/or its predisposition for meat marbling.

Several documents are cited throughout the text of this specification. The disclosure content of the documents cited herein (including any manufacture's specifications, instructions, etc.) is herewith incorporated by reference.

Milk fat content is a continuously distributed trait with heritability estimates between 0.45 and 0.50 (Goddard and Wiggans, 1999). There are considerable differences in the average milk fat content between different cattle breeds, ranging from 3.6% in the Holstein to 4.6% in the Jersey breed. The systematic mapping of quantitative trait loci (QTL) underlying the genetic variance of milk production traits resulted in approximate map positions of QTL for milk fat content (Georges *et al.*, 1995; Zhang *et al.*, 1998; Heyen *et al.*, 1999; Velmala *et al.*, 1999). The most consistent results were reported for a QTL on chromosome 14 (Coppieters *et al.*, 1998) (Riquet *et al.*, 1999). The mapping interval of this QTL could be reduced to a few Centimorgans. High-resolution comparative maps of the critical region did not reveal obvious positional candidate genes (Riquet *et al.*, 1999). *DGAT*, the gene encoding acyl CoA:diacylglycerol transferase, a microsomal enzyme that catalyses the final step of triglyceride synthesis, became a functional candidate after it had been shown that mice lacking both copies of *DGAT* show defective lactation. This is most likely the consequence of deficient triglyceride synthesis in the mammary gland (Smith *et al.*, 2000).

Another candidate was reported by Barendse *et al.* (1999). They described a polymorphism in the 5' untranslated region of the gene encoding thyroglobulin (*TG*) which was postulated to be associated with lipid metabolism, particularly the

deposition of fat in muscular tissue. Said deposition of fat produces the typical marbling of the meat. The gene was localized on bovine chromosome 14 very close to the *DGAT* locus (Threadgill et al. 1990). However, the protein encoded by the gene *TG* is not involved in triglyceride synthesis and thus fat deposition.

In summary, the state of the art did so far not provide any genetic link with fat content in milk that can be efficiently used in routine testing.

Thus and in of the above, the technical problem underlying the present invention was to provide a method of testing mammals for their predisposition for fat content of milk and/or its predisposition for meat marbling. Said method ought to be easy to use and offer the opportunity to conveniently analyze large numbers of samples. The solution to this technical problem is achieved by providing the embodiments characterized in the claims.

Accordingly the present invention relates to a nucleic acid molecule encoding a bovine acyl CoA:diacylglycerol transferase (*DGAT*) contributing to or indicative for low fat content of milk and to low meat marbling (intramuscular fat content); wherein said nucleic acid molecule is selected from the group consisting of:

- (a) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO: 1;
- (b) a nucleic acid molecule comprising the coding sequence of the polypeptide of SEQ ID NO: 2;
- (c) a nucleic acid molecule the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (a) or (b), wherein said nucleic acid molecule has at the position corresponding to position 10433 and 10434 of the *DGAT* gene (SEQ ID NO: 1) a guanine and a cytosine residue; and
- (d) a nucleic acid molecule the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (a) or (b), wherein said nucleic acid molecule has at the *DGAT* gene (SEQ ID NO: 1) position
 - (i) ... 3343 a cytosine, 10433 a guanine, 10434 a cytosine, 11030 a guanine, 11048 a cytosine and 11093 a thymine;

- (ii) 3343 a cytosine, 10433 a guanine, 10434 a cytosine, 11030 a guanine, 11048 a thymine, and 11093 a thymine; or
- (iii) 3343 a guanine, 10433 a guanine, 10434 a cytosine, 11030 a guanine, 11048 a thymine and 11093 a thymine.

Genetic screening (also called genotyping or molecular screening), can be broadly defined as testing to determine if an individual has mutations (alleles or polymorphisms) that either cause a specific phenotype or are "linked" to the mutation causing the phenotype. Linkage refers to the phenomenon that the DNA sequences which are close together in the genome have a tendency to be inherited together. Two or more sequences may be linked because of some selective advantage of co-inheritance. More typically, however, two or more polymorphic sequences are co-inherited because of the relative infrequency with which meiotic recombination events occur within the region between the two polymorphisms. The co-inherited polymorphic alleles are said to be in linkage disequilibrium with one another because, in a given population, they tend to either both occur together or else not occur at all in any particular member of the population. Indeed, where multiple polymorphisms in a given chromosomal region are found to be in linkage disequilibrium with one another, they define a quasi-stable genetic "haplotype."

Furthermore, where a phenotype-causing mutation is found within or in linkage with this haplotype, one or more polymorphic alleles of the haplotype can be used as a diagnostic or prognostic indicator of the likelihood of developing a specific phenotype. Identification of a haplotype which spans or is linked to a phenotype-causing mutational change, serves as a predictive measure of an individual's likelihood of having inherited that phenotype-causing mutation. Importantly, such prognostic or diagnostic procedures can be utilized without necessitating the identification and isolation of the actual phenotype-causing molecule. This is significant because the precise determination of the molecular basis of the establishment of a specific phenotype can be difficult and laborious, especially in the case of multifactorial phenotype.

Mapping studies on human chromosome 8 placed *DGAT* indirectly within the mapping interval of the QTL on bovine chromosome 14, the homologous

counterpart of human chromosome 8. Sequencing of *DGAT* from pooled DNA revealed massive frequency shifts at several variable positions between groups of animals with high and low milk fat percentage, respectively. The procedure of said sequencing is described in example 6. It was searched for variation in 10528 basepairs, i.e., the entire coding region of *DGAT*, the major part of the introns and the 5' and 3' regions. 20 variable positions were identified, mostly single nucleotide polymorphisms (summarized in table 9). By said method several nucleotide polymorphisms were detected which were unexpected vis-à-vis the prior art data for the sequences known from the region the *DGAT* in mice, human or plants. Among the variants is a double substitution causing the non-conservative substitution of alanine by lysine. Furthermore, said variants comprised several single nucleotide substitutions. An example for a sequence containing said newly identified polymorphisms is SEQ ID NO: 1.

Direct sequencing in animals belonging to different breeds of *Bos taurus taurus* and *Bos taurus indicus* as well as in animals of *Bos grunniens* (yak) and *Bubalus bubalus* (water buffalo) at position 3343, 10433, 10434, 11030, 11048 and 11093 allowed to derive at least 8 haplotypes (see Fig. 12). The haplotypes observed encoded a DGAT1 protein with either a lysine or an alanine in position 232 of the DGAT1 polypeptide sequence. In addition, specific nucleotides at positions 3343, 10433, 10434, 11030, 11048 and 11093 were demonstrated to be indicative of a specific haplotype. As shown in Fig. 12A, haplotypes encoding a protein with a lysine in position 232 may contain in the above mentioned positions either TAAGCC, CAAGCC, CAAGCT, CAAACC or CAAACT while alanine encoding haplotypes are characterized by CGCGCT (i.e. at position: 3343 cytosine, 10433 a guanine, 10434 a cytosine, 11030 a guanine, 11048 a cytosine and 11093 a thymine), CGCGTT (i.e. at position: 3343 a cytosine, 10433 a guanine, 10434 a cytosine, 11030 a guanine, 11048 a thymine, and 11093 a thymine) or GGCGTT (i.e. at position: 3343 a guanine, 10433 a guanine, 10434 a cytosine, 11030 a guanine, 11048 a thymine and 11093 a thymine) in the above mentioned positions. It is of note that the invention also comprises sequences wherein one or two nucleotides in the above-indicated positions are exchanged by different nucleotides. In addition, the invention comprises haplotypes arising from recombination events and including the above recited gene.

Furthermore, an RFLP analysis revealed frequency estimates for lysine and alanine encoding alleles in several cattle breeds of Bovinae subfamilies (see Fig. 12b). Distinct frequency differences for the allelic distribution in various breeds indicated a correlation between milk fat content and the genetic variation.

The term "hybridizes under stringent conditions", as used in the description of the present invention, is well known to the skilled artisan and corresponds to conditions of high stringency. Appropriate stringent hybridization conditions for each sequence may be established by a person skilled in the art on well-known parameters such as temperature, composition of the nucleic acid molecules, salt conditions etc.; see, for example, Sambrook et al., "Molecular Cloning, A Laboratory Manual"; CSH Press, Cold Spring Harbor, 1989 or Higgins and Hames (eds.), "Nucleic acid hybridization, a practical approach", IRL Press, Oxford 1985, see in particular the chapter "Hybridization Strategy" by Britten & Davidson, 3 to 15. Stringent hybridization conditions are, for example, conditions comprising overnight incubation at 42° C in a solution comprising: 50% formamide, 5x SSC (750 mM NaCl, 75 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°. Other stringent hybridization conditions are for example 0.2 x SSC (0.03 M NaCl, 0.003M Natriumcitrat, pH 7) bei 65°C. Preferred in accordance with the present invention are nucleic acids which are capable of hybridizing to the nucleic acid molecule of the invention or parts thereof wherein said nucleic acid molecule has at the position corresponding to position 10433 and 10434 of the *DGAT* gene (SEQ ID NO: 1) a guanine and a cytosine residue. More preferred in accordance with the present invention are nucleic acids which are capable of hybridizing to the complementary strand of any of the nucleic acid molecules of the invention or parts thereof, wherein said nucleic acid molecule contains at position 3343, 10433, 10434, 11030, 11048 and 11093 of the *DGAT* gene (SEQ ID NO: 1) nucleotides which are either CGCGCT, CGCGTT or GGCGTT. Furthermore, the nucleic acid molecules of the invention may contain any alanine codon at the position encoding amino acid 232 of *DGAT*.

The term "corresponding" as used herein means that a position is not only determined by the number of the preceding nucleotides and amino acids,

respectively. The position of a given nucleotide or amino acid in accordance with the present invention which may be deleted, substituted or comprise one or more additional nucleotide(s) may vary due to deletions or additional nucleotides or amino acids elsewhere in the gene or the polypeptide. Thus, under a "corresponding position" in accordance with the present invention it is to be understood that nucleotides or amino acids may differ in the indicated number but may still have similar neighboring nucleotides or amino acids. Said nucleotides or amino acids may for instance together with their neighbors form sequences which may be involved in the regulation of gene expression, stability of the corresponding RNA or RNA editing, as well as encode functional domains or motifs of the protein of the invention. In the context of the invention functional domains or motifs of the invention are defined as portions having the enzymatic activity of *DGAT* and/or portions which are capable to be recognized as an antigen and therefore represent an epitope for an antibody or small molecule.

Therefore, the invention comprises allelic variants of the *DGAT* gene as well as recombinantly or otherwise altered *DGAT* sequences. In conformance with the present invention, the recited nucleic acid "encodes" the *DGAT* enzyme. Whereas by definition the claimed nucleic acid molecule comprises the coding region, it may also comprise non-coding regions such as regulatory regions or introns.

Apart from being the subject of investigation, the nucleic acid molecule of the invention may be useful as probes in Northern or Southern Blot analysis of RNA or DNA preparations, respectively, or can be used as oligonucleotide primers in PCR analysis dependent on their respective size. Also comprised by the invention are hybridizing nucleic acids which are useful for analyzing DNA-Protein interactions via, e.g., electrophoretic mobility shift analysis (EMSA). Preferably, said hybridizing nucleic acids comprise at least 10, more preferably at least 15 nucleotides in length while a hybridizing polynucleotide of the present invention to be used as a probe preferably comprises at least 100, more preferably at least 200, or most preferably at least 500 nucleotides in length.

The nucleic acid molecule of the invention is expected to occur in any breed of the bovine species. In a preferred embodiment of the invention the bovine nucleic acid molecule is a nucleic acid molecule of a bovine animal selected from the group

consisting of Ayrshire, Bazadaise, Beefalo, Blaarkop, Braunvieh Fleischnutzung, Grauvieh, Lakenfelder, Limpurger Fleischnutzung, Maine Anjou, Marchigiana, Montbeliard, Murnau-Werdenfelser, Normanne, Romagnola, Rotbunt Fleischnutzung, Telemark, Tuxer, Vogesen-Rind, Wasserbüffel, Witrug, Yak, Auerochse, Bison/Wisent, Hinterwälder Fleischnutzung, Vorderwälder Fleischnutzung, Angler, Doppelnutzung Rotbunt, Holstein-Rbt., Holstein-Sbt., Holstein-Friesian, Deutsches Shorthorn, Rotvieh alter Angler, Aberdeen Angus, Aubrac, Blonde d'Aquitaine, Brahman, Brangus, Charolais, Chlanina, Deutsche Angus, Fjall-Rind, Fleckvieh Fleischnutzung Ost, Gelbvieh Fleischnutzung, Hereford, Jersey, Limousin, Lincoln Red, Piemonteser, Salers, South Devon, Weißblaue Belgier, Beited Galloway, Dexter, Galloway, Highland, Longhorn, Luing, Ungarisches Steppenrind, Welsh-Black, White Galloway, White Park, Zwerg-Zebus, Rotvieh Zuchtrichtung, Uckermärker, Deutsche Schwarzbunte alter, Braunvieh, Fleckvieh, Gelbvieh, Pinzgauer Fleischnutzung, Ansbach-Triesdorfer, Braunvieh alter Zuchtrichtung, Limpurger, Murnau-Werdenfelser, Pinzgauer, Pustertaler Schecken, Hinterwäldler, Vorderwäldler and Glanrind.

In a more preferred embodiment of the invention the bovine nucleic acid molecule is a nucleic acid molecule of a female bovine animal.

The nucleic acid molecule can be taken from any nucleic acid containing tissue. Preferably said nucleic acid molecule is present in a sample taken from, for example, from muscle, blood, skin, milk, urine and other samples taken from a bovine animal.

Preferably said nucleic acid molecule is mRNA, genomic DNA (gDNA) or cDNA which is derived from said mRNA by reverse transcription of said mRNA.

The method of reverse transcription of mRNA into cDNA is well established and known by a person skilled in the art.

More preferably said gDNA is a gene.

In an preferred embodiment of the invention the nucleic acid molecule is a fragment of the herein above described nucleic acid molecule having at least 14 nucleotides wherein said fragment comprises nucleotide position 10433 and 10434 of SEQ ID NO: 1.

Said nucleic acid molecule may, for example, be used as hybridization probe. For hybridization probes, it may be, e.g., desirable to use nucleic acid analogs, in order to improve the stability and binding affinity. The term "nucleic acid" shall be understood to encompass such analogs. A number of modifications have been described that alter the chemistry of the phosphodiester backbone, sugars or heterocyclic bases. Among useful changes in the backbone chemistry are phosphorothioates; phosphorodithioates, where both of the non-bridging oxygens are substituted with sulfur; phosphoroamidites; alkyl phosphotriesters and boranophosphates. Achiral phosphate derivatives include 3'-O'-5'-S-phosphorothioate, 3'-S-5'-O-phosphorothioate, 3'-CH₂-5'-O-phosphonate and 3'-NH-5'-O-phosphoroamidate. Peptide nucleic acids replace the entire phosphodiester backbone with a peptide linkage. Sugar modifications are also used to enhance stability and affinity. The α -anomer of deoxyribose may be used, where the base is inverted with respect to the natural β -anomer. The 2'-OH of the ribose sugar may be altered to form 2'-O-methyl or 2'-O-allyl sugars, which provides resistance to degradation without comprising affinity. Modification of the heterocyclic bases must maintain proper base pairing. Some useful substitutions include deoxyuridine for deoxythymidine; 5-methyl-2'-deoxycytidine and 5-bromo-2'-deoxycytidine for deoxycytidine. 5-propynyl-2'-deoxyuridine and 5-propynyl-2'-deoxycytidine have been shown to increase affinity and biological activity when substituted for deoxythymidine and deoxycytidine, respectively.

The hybridization probe or the primer(s) used for amplification may also contain a detectable label. Suitable labels include fluorochromes, e.g. fluorescein isothiocyanate (FITC), rhodamine, Texas Red, phycoerythrin, allophycocyanin, 6-carboxyfluorescein (6-FAM), 2',7'-dimethoxy-4',5'-dichloro-6-carboxyfluorescein (JOE), 6-carboxy-X-rhodamine(ROX), 6-carboxy-2',4',7',4,7-hexachlorofluorescein (HEX), 5-carboxyfluorescein (5-FAM) or N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA), radioactive labels, e.g. ³²P, ³⁵S, ³H; etc. The label may also be a two stage system, where the DNA is conjugated to biotin, haptens, etc. having a high

affinity binding partner, e.g. avidin, specific antibodies, etc., where the binding partner is conjugated to a detectable label. In the case of amplification the label may be conjugated to one or both of the primers. The pool of nucleotides used in the amplification may also be labeled, so as to incorporate the label into the amplification product. Alternatively, the double strand formed after hybridization can be detected by anti-double strand DNA specific antibodies or aptamers etc.

More preferably said nucleic acid molecule is complementary to the above described nucleic acid. Said complementary nucleic acid molecule is suitable to hybridize specifically with a polynucleotide as described above. Specific hybridization occurs preferably under stringent conditions and implies no or very little cross-hybridization with nucleotide sequences encoding no or substantially different proteins. Such nucleic acid molecules may be used as probes and/or for the control of gene expression. Nucleic acid probe technology is well known to those skilled in the art who will readily appreciate that such probes may vary in length. Preferred are nucleic acid probes of 17 to 35 nucleotides in length. Of course, it may also be appropriate to use nucleic acids of up to 100 and more nucleotides in length. The nucleic acid probes of the invention are useful for various applications. On the one hand, they may be used as PCR primers for amplification of nucleic acid molecules according to the invention. Another application is the use as a hybridization probe to identify polynucleotides hybridizing to the nucleic acid molecule of the invention by homology screening of genomic DNA libraries (see example 3). Nucleic acid molecules according to this preferred embodiment of the invention which are complementary to a polynucleotide as described above may also be used for repression of expression of a gene comprising such a polynucleotide, for example due to an antisense or triple helix effect or for the construction of appropriate ribozymes (see, e.g., EP-A1 0 291 533, EP-A1 0 321 201, EP-A2 0 360 257) which specifically cleave the (pre)-mRNA of a gene comprising a polynucleotide of the invention. Selection of appropriate target sites and corresponding ribozymes can be done as described for example in Steinecke, Ribozymes, Methods in Cell Biology 50, Galbraith et al. eds Academic Press, Inc. (1995), 449-460. Standard methods relating to antisense technology have also been described (Melani, Cancer Res. 51 (1991), 2897-2901). Furthermore, the

person skilled in the art is well aware that it is also possible to label such a nucleic acid probe with an appropriate marker for specific applications, such as for the detection of the presence of a polynucleotide of the invention in a sample derived from an organism.

The above described nucleic acid molecules may either be DNA or RNA or a hybrid thereof. Furthermore, said nucleic acid molecule may contain, for example, thioester bonds and/or nucleotide analogues, commonly used in oligonucleotide anti-sense approaches. Said modifications may be useful for the stabilization of the nucleic acid molecule against endo- and/or exonucleases in the cell. Said nucleic acid molecules may be transcribed by an appropriate vector containing a chimeric gene which allows for the transcription of said nucleic acid molecule in the cell. Such nucleic acid molecules may further contain ribozyme sequences as described above.

Furthermore, the present invention provides a vector comprising the herein above described nucleic acid molecule. Said expression vectors may particularly be plasmids, cosmids, viruses or bacteriophages used conventionally in genetic engineering plasmids, cosmids, viruses and bacteriophages used conventionally in genetic engineering that comprise the aforementioned nucleic acid. Preferably, said vector is a gene transfer or targeting vector. Expression vectors derived from viruses such as retroviruses, vaccinia virus, adeno-associated virus, herpes viruses, or bovine papilloma virus, may be used for delivery of the nucleic acid into targeted cell population. Methods which are well known to those skilled in the art can be used to construct recombinant viral vectors; see, for example, the techniques described in Sambrook et al., *Molecular Cloning A Laboratory Manual*, Cold Spring Harbor Laboratory (1989) N.Y. and Ausubel et al., *Current Protocols in Molecular Biology*, Green Publishing Associates and Wiley Interscience, N.Y. (1989). Alternatively, the nucleic acids and vectors can be reconstituted into liposomes for delivery to target cells. The vectors containing the nucleic acid can be transferred into the host cell by well-known methods, which vary depending on the type of cellular host. For example, calcium phosphate or DEAE-Dextran mediated transfection or electroporation may be used for eukaryotic cellular hosts; see Sambrook, supra. Such vectors may comprise further genes such as marker genes

which allow for the selection of said vector in a suitable host cell and under suitable conditions.

Preferably, said vector comprises regulatory elements for expression of said nucleic acid molecule. Consequently, the nucleic acid of the invention may be operatively linked to expression control sequences allowing expression in eukaryotic cells. Expression of said nucleic acid molecule comprises transcription of the sequence nucleic acid molecule into a translatable mRNA. Regulatory elements ensuring expression in eukaryotic cells, preferably mammalian cells, are well known to those skilled in the art. They usually comprise regulatory sequences ensuring initiation of transcription and, optionally, a poly-A signal ensuring termination of transcription and stabilization of the transcript, and/or an intron further enhancing expression of said nucleic acid. Additional regulatory elements may include transcriptional as well as translational enhancers, and/or naturally-associated or heterologous promoter regions. Possible regulatory elements permitting expression in eukaryotic host cells are the AOX1 or GAL1 promoter in yeast or the CMV-, SV40-, RSV-promoter (Rous sarcoma virus), CMV-enhancer, SV40-enhancer or a globin intron in mammalian and other animal cells. Beside elements which are responsible for the initiation of transcription such regulatory elements may also comprise transcription termination signals, such as the SV40-poly-A site or the tk-poly-A site, downstream of the nucleic acid molecule. Furthermore, depending on the expression system used leader sequences capable of directing the polypeptide to a cellular compartment or secreting it into the medium may be added to the coding sequence of the aforementioned nucleic acid and are well known in the art. The leader sequence(s) is (are) assembled in appropriate phase with translation, initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein, or a portion thereof, into the periplasmic space or extracellular medium. Optionally, the heterologous sequence can encode a fusion protein including an C- or N-terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product. In this context, suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDVI (Pharmacia), pCDM8, pRc/CMV, pcDNA1, pcDNA3,

the Echo™ Cloning System (Invitrogen), pSPORT1 (GIBCO BRL) or pRevTet-On/pRevTet-Off or pCI (Promega).

Another preferred embodiment of the invention relates to primer or primer pair, wherein the primer or primer pair hybridize under stringent conditions to the nucleic acid molecule of the invention comprising nucleotide position 10433 and 10434 of SEQ ID NO: 1 or the complement strand thereof. The exact composition of the primer sequences is not critical as long as they allow detection of the desired sequence(s). Preferably, the primers are chosen in such a way that they hybridize under stringent conditions to the desired sequence(s). It is preferable to choose a primer or a pair of primers that will generate an amplification product of at least 50 nt, preferably of at least about 100 nt and most preferably of at least 200 nt. Algorithms for the selection of primer sequences are generally known and are available in commercial software packages (see example 1). Amplification primers hybridize to complementary strands of DNA and will prime towards each other.

Furthermore, the present invention relates to a host cell which contains the herewith above described expression vector.

Preferably, said host cell is a eukaryotic, most preferably a mammalian cell if therapeutic uses of the protein are envisaged. Of course, yeast and less preferred prokaryotic, e.g., bacterial cells may serve as well, in particular if the produced protein is used as a diagnostic means.

The polynucleotide or vector of the invention which is present in the host cell may either be integrated into the genome of the host cell or it may be maintained extrachromosomally.

The term "prokaryotic" is meant to include all bacteria which can be transformed or transfected with a DNA or RNA molecules for the expression of a protein of the invention. Prokaryotic hosts may include gram negative as well as gram positive bacteria such as, for example, E. coli, S. typhimurium, Serratia marcescens and Bacillus subtilis. The term "eukaryotic" is meant to include yeast, higher plant, insect and preferably mammalian cells. Depending upon the host employed in a recombinant production procedure, the protein encoded by the polynucleotide of the present invention may be glycosylated or may be non-glycosylated. A nucleic acid

molecule of the invention can be used to transform or transfect the host using any of the techniques commonly known to those of ordinary skill in the art. Furthermore, methods for preparing fused, operably linked genes and expressing them in, e.g., mammalian cells and bacteria are well-known in the art (Sambrook, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1989). The genetic constructs and methods described therein can be utilized for expression of the protein of SEQ ID NO: 2 in eukaryotic or prokaryotic hosts. In general, expression vectors containing promoter sequences which facilitate the efficient transcription of the inserted polynucleotide are used in connection with the host. The expression vector typically contains an origin of replication, a promoter, and a terminator, as well as specific genes which are capable of providing phenotypic selection of the transformed cells.

In an alternative embodiment the present invention relates to a method for production of a functional bovine *DGAT* or a functional fragment thereof comprising:

- (a) culturing said host cell containing the expression vector which comprises the herein above mentioned nucleic acid molecule under conditions allowing the expression of the encoded polypeptide; and
- (b) collecting the polypeptide from the culture.

As aforementioned, a functional fragment is defined in the context of the present invention as a fragment having the enzymatic activity of *DGAT* and/or fragment which is capable to be recognized as an antigen and therefore represent an epitope for an antibody and/or small molecule suitable for specific binding and detection of an epitope.

The transformed hosts can be grown in fermentors and cultured according to techniques known in the art to achieve optimal cell growth. The protein of the invention can then be isolated from the growth medium, cellular lysates, or cellular membrane fractions. Once expressed, the protein of the present invention can be purified according to standard procedures of the art, including ammonium sulfate precipitation, affinity columns, column chromatography, gel electrophoresis and the like; see, Scopes, "Protein Purification", Springer-Verlag, N.Y. (1982). Substantially pure proteins of at least about 90 to 95% homogeneity are preferred, and 98 to 99% or more homogeneity are most preferred, for pharmaceutical uses. Once purified,

partially or to homogeneity as desired, the proteins may then be used therapeutically (including extracorporeally) or in developing and performing assay procedures.

Hence, in a still further embodiment, the present invention relates to functional bovine *DGAT* polypeptide as depicted in SEQ ID NO: 2 or a functional fragment thereof encoded by a nucleic acid molecule (SEQ ID NO: 1) or produced by a method of as described above. It will be apparent to those skilled in the art that the protein of the invention can be further coupled to other moieties for, e.g., drug targeting and imaging applications. Such coupling may be conducted chemically after expression of the protein to site of attachment or the coupling product may be engineered into the protein of the invention at the DNA level. The DNAs are then expressed in a suitable host system, and the expressed proteins are collected and renatured, if necessary.

Furthermore, the provision of the protein of the present invention enables the production of *DGAT* specific antibody which binds to an epitope of the polypeptide or fragment of SEQ ID NO: 2 the epitope comprising a alanine at position 232 but not to a polypeptide or a fragment of SEQ ID NO: 4 having a lysine at position 232. In an alternative embodiment the invention relates to the production of *DGAT* specific antibody which binds to an epitope of the polypeptide or fragment of SEQ ID NO: 4 the epitope comprising a lysine at position 232 but not to a polypeptide or a fragment of SEQ ID NO: 2 having a alanine at position 232.

In this respect, hybridoma technology enables production of cell lines secreting antibody to essentially any desired substance that produces an immune response. RNA encoding the light and heavy chains of the immunoglobulin can then be obtained from the cytoplasm of the hybridoma. The 5' end portion of the mRNA can be used to prepare cDNA to be inserted into an expression vector. The DNA encoding the antibody or its immunoglobulin chains can subsequently be expressed in cells, preferably mammalian cells.

Depending on the host cell, renaturation techniques may be required to attain proper conformation of the antibody. If necessary, point substitutions seeking to

optimize binding may be made in the DNA using conventional cassette mutagenesis or other protein engineering methodology such as is disclosed herein.

Said antibodies, which are monoclonal antibodies, polyclonal antibodies, single chain antibodies, or fragment thereof that specifically binds said peptide or polypeptide also including bispecific antibody, synthetic antibody, antibody fragment, such as Fab, a F(ab₂)', Fv or scFv fragments etc., or a chemically modified derivative of any of these (all comprised by the term "antibody"). Monoclonal antibodies can be prepared, for example, by the techniques as originally described in Köhler and Milstein, Nature 256 (1975), 495, and Galfré, Meth. Enzymol. 73 (1981), 3, which comprise the fusion of mouse myeloma cells to spleen cells derived from immunized mammals with modifications developed by the art. Furthermore, antibodies or fragments thereof to the aforementioned peptides can be obtained by using methods which are described, e.g., in Harlow and Lane "Antibodies, A Laboratory Manual", CSH Press, Cold Spring Harbor, 1988. When derivatives of said antibodies are obtained by the phage display technique, surface plasmon resonance as employed in the BIAcore system can be used to increase the efficiency of phage antibodies which bind to an epitope of the peptide or polypeptide of the invention (Schier, Human Antibodies Hybridomas 7 (1996), 97-105; Malmberg, J. Immunol. Methods 183 (1995), 7-13). The production of chimeric antibodies is described, for example, in WO89/09622. A further source of antibodies to be utilized in accordance with the present invention are so-called xenogenic antibodies. The general principle for the production of xenogenic antibodies such as human antibodies in mice is described in, e.g., WO 91/10741, WO 94/02602, WO 96/34096 and WO 96/33735. Antibodies to be employed in accordance with the invention or their corresponding immunoglobulin chain(s) can be further modified using conventional techniques known in the art, for example, by using amino acid deletion(s), insertion(s), substitution(s), addition(s), and/or recombination(s) and/or any other modification(s) known in the art either alone or in combination. Methods for introducing such modifications in the DNA sequence underlying the amino acid sequence of an immunoglobulin chain are well known to the person skilled in the art; see, e.g., Sambrook, Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory (1989) N.Y.

Moreover, the present invention relates to a transgenic, non-human animal comprising at least the herein above disclosed nucleic acid molecules. Preferably said transgenic, non-human animal belongs to cattle.

In an other embodiment the present invention relates to a method of testing a mammal for its predisposition for fat content of milk and/or its predisposition for meat marbling comprising analyzing the nucleic acid of a sample comprising the gene encoding *DGAT*, a corresponding mRNA for nucleotide polymorphisms which are connected with said predisposition or any nucleic acid molecule of the invention. The term "its predisposition for fat content of milk and/or its predisposition for meat marbling" describes the capability of a mammal to produce milk with high fat, respectively low fat content and/or its capability to produce meat with high intramuscular fat content, respectively low intramuscular fat content.

Preferably the nucleic acid of said method is DNA.

More preferably the nucleic acid of said method is gDNA (genomic DNA).

Also more preferred the nucleic acid is cDNA which is derived from said mRNA by reverse transcription of said mRNA.

In accordance with the invention the nucleotide polymorphisms which are contributing to or indicative for low fat content of milk and to low meat marbling are in one preferred embodiment located in the coding region of the *DGAT* gene.

More preferably the nucleotide polymorphisms in the coding region of the gene encoding *DGAT* result in substitution, deletion and/or addition of at least one amino acid in the amino acid sequence of the polypeptide which is encoded by said gene.

Further more preferably said nucleic acid molecule has at the position corresponding to position 10433 and 10434 of the *DGAT* gene (SEQ ID NO: 1) a guanine and a cytosine residue which corresponds to i.e. correlates with a predisposition for low fat content of milk and low meat marbling.

More preferably the nucleic acid molecule has at the positions corresponding to position 3343, 10433, 10434, 11030, 11048 and 11093 of the *DGAT* gene (SEQ ID NO:1) the nucleotides CGCGCT (i.e. at position 3343 a cytosine, 10433 a guanine, 10434 a cytosine, 11030 a guanine, 11048 a cytosine and 11093 a thymine), CGCGTT (i.e. at position 3343 a cytosine, 10433 a guanine, 10434 a cytosine, 11030 a guanine, 11048 a thymine, and 11093 a thymine) or GGCGTT (i.e. at position 3343 a guanine, 10433 a guanine, 10434 a cytosine, 11030 a guanine, 11048 a thymine and 11093 a thymine) which corresponds to i.e. correlates with a predisposition for low fat content of milk and low meat marbling.

Alternatively said nucleic acid molecule has at the position corresponding to position 10433 and 10434 of the *DGAT* gene (SEQ ID NO: 3) two adenine residue which corresponds to i.e. correlates with a predisposition for high fat content of milk and high meat marbling.

More preferably said nucleic acid molecule has at the positions corresponding to positions 3343, 10433, 10434, 11030, 11048 and 11093 of the *DGAT* gene the nucleotides TAAGCC (i.e. at position 3343 a thymine, 10433 an adenosine, 10434 an adenosine, 11030 a guanine, 11048 a cytosine and 11093 a cytosine), CAAGCC (i.e. at position 3343 a cytosine, 10433 an adenosine, 10434 an adenosine, 11030 a guanine, 11048 a cytosine, and 11093 a cytosine), CAAGCT (i.e. at position 3343 a cytosine, 10433 an adenosine, 10434 an adenosine, 11030 a guanine, 11048 a cytosine and 11093 a thymine), CAAACC (i.e. at position 3343 a cytosine, 10433 an adenosine, 10434 an adenosine, 11030 an adenosine, 11048 a cytosine and 11093 a cytosine) or CAAACT (i.e. at position 3343 a cytosine, 10433 an adenosine, 10434 an adenosine, 11030 an adenosine, 11048 a cytosine and 11093 a thymine) which corresponds to i.e. correlates with a predisposition for high fat content of milk and high meat marbling.

Also in accordance with the invention the nucleotide polymorphisms are preferably located in a region which is responsible for the regulation of the expression of the product of the gene encoding *DGAT*.

More preferred the nucleotide polymorphisms which are analyzed by the method of the invention are single nucleotide polymorphisms (SNP).

In another preferred embodiment said testing in the method of the invention comprises hybridizing a herein above described nucleic acid molecule as a probe under stringent conditions to the nucleic acid molecules comprised in said sample and detecting hybridization. Such stringent conditions are known by a person skilled in the art and also described herein above.

More preferably said testing comprises digesting the product of said hybridization with a restriction endonuclease and analyzing the product of said digestion.

Even more preferred said probe is detectably labeled.

Alternatively, said testing comprises determining the nucleic acid sequence of at least a portion of said nucleic acid molecule. Methods for sequencing of nucleic acids are known in the art. An example for said testing for predisposition of individual animals by comparative sequencing is described herein below in example 6.

Preferably said determination of the nucleic acid sequence is effected by solid-phase minisequencing.

Also alternatively the testing further comprises, prior to analyzing the nucleic acid, amplification of at least a portion of said nucleic acid.

More preferred in said amplification reaction at least one of the primers employed in said amplification reaction is the primer or belongs to the primer pair as aforementioned, the method comprising assaying for an amplification product.

Even more preferred said amplification is effected by or said amplification is the polymerase chain reaction (PCR).

Furthermore, alternatively the method of the invention further comprises analyzing said nucleic acid by the use of:

- (a) a primer extension assay;
- (b) a differential hybridization assay; and/or
- (c) an assay which detects allele-specific enzyme cleavage.

The underlying principles and the use of said assays has been described in an article of Asil Memisoglu (www.thebiotechclub.org/Tech/pharmacogenomics.html). Examples for said assays are known by a person skilled in the art. Furthermore, the method of analyzing said nucleic acid by the use of an assay which detects allele-specific enzyme cleavage is describe in example 8 herein below.

Furthermore, in an other embodiment the invention relates to a method of testing a mammal for its predisposition for fat content of milk and/or its predisposition for meat marbling, said method comprising the steps of:

- (a) preparation of a tissue sample from the subject;
- (b) contacting the sample with an aforementioned antibody specifically binding to an epitope of the polypeptide or fragment of SEQ ID NO: 2 the epitope comprising a alanine at position 232 but not to a polypeptide or a fragment of SEQ ID NO: 4 having a lysine at position 232 or specifically binding to an epitope of the polypeptide or fragment of SEQ ID NO: 4 the epitope comprising a lysine at position 232 but not to a polypeptide or a fragment of SEQ ID NO: 2 having a alanine at position 232; and
- (c) detecting whether a specific binding of said antibody to its antigen has occurred.

Said method may comprise the transfer of the sample onto a membrane, e.g. by blot technique after electrophoresis. If so the detection whether a specific binding has occurred may comprise washing of the membrane to remove agent unspecifically bound to the membrane. Said detection may be performed by the use of agents which on the one hand are suitable for the detection of the presence of the specifically interacting agent. Furthermore said agents may comprises a domain or function which can be used for the generation of a detectable signal. The steps of contacting the proteins with said agents and detecting whether a specific interaction

has occurred may be similar to the principle of immunodetection of proteins by Western Blot known to the person skilled in the art.

Preferably said method wherein the binding of the antibody which specifically binds to an epitope of the polypeptide or fragment of SEQ ID NO: 2 the epitope comprising a alanine at position 232 but not to a polypeptide or a fragment of SEQ ID NO: 4 having a lysine at position 232 indicates a predisposition of the mammal for low fat content of milk and to low meat marbling.

Also preferred, said method wherein the binding of the antibody which specifically binds to an epitope of the polypeptide or fragment of SEQ ID NO: 4 the epitope comprising a lysine at position 232 but not to a polypeptide or a fragment of SEQ ID NO: 2 having a alanine at position 232 indicates a predisposition of the mammal for high fat content of milk and to high meat marbling.

Also preferred is a method for testing of a mammal for its predisposition for low fat content and/or its predisposition for meat marbling comprising analyzing nucleotide positions 3343, 10433, 10434, 11030, 11048 and 11093 of the DGAT gene (SEQ ID NO:1), wherein the nucleotides CGCGCT, CGCGTT or GGCGTT at the above-indicated positions are indicative of low fat content of milk and low meat marbling.

Also preferred is a method for testing of a mammal for its predisposition for high fat content and/or its predisposition for meat marbling comprising analyzing nucleotide positions 3343, 10433, 10434, 11030, 11048 and 11093 of the DGAT gene (SEQ ID NO:1), wherein the nucleotides TAAGCC, CAAGCC, CAAGCT, CAAACC or CAAACT at the above-indicated positions are indicative of high fat content of milk and high meat marbling.

More preferred the samples which are analyzed by the methods of the invention are isolated from cloven hoofed animals.

In a further more preferred embodiment said cloven hoofed animals are cattle, buffalos, yaks or pigs.

Finally the present invention relates in one embodiment to a kit comprising at least the aforementioned fragment, the aforementioned nucleic acid molecule, the aforementioned primer or primer pair , or one of the aforementioned in one or more containers.

The figures show

Figure 1 Bovine metaphase spread after fluorescence in situ hybridization using BAC clone 56-F1. BAC-DNA was labeled with biotin using nick-translation. Detection of the hybridized probe was performed with streptavidin-Cy3. Photos were taken with a CCD-camera coupled to a Zeiss microscope with a magnification of 650 x. The signals on both copies of chromosome 14 are indicated by arrow and arrow head. Note that one copy of chromosome 14 (signal indicated by arrow) is involved in a Robertsonian fusion with chromosome 20.

Figure 2 Partial maps of three BACs (56-F1, 240-A1, 269-H17). Solid lines represent sequenced parts. The vector sequences are shown as gray boxes. T7 and SP6 refer to the primers used for BAC-end sequencing. The colored boxes represent genes: *DGAT*, diacylglycerol acyltransferase; *HSF1*, heat shock transcription factor 1; *FPXL6*, f-box and leucine-rich repeat protein 6. Annotation of the sequences is based on a high similarity with the corresponding human sequences. The arrows indicate the orientation of the genes. Drawings are not to scale.

Figure 3 EST-derived transcript map of the bovine *DGAT* gene. The blue areas represent sequences covered by the ESTs. T0 is composed of ESTs AW483961, AW486026, AW652329, BE664362, BE753833, BE664357, T1 of AW446908, T2 of AW446985, T4 of AW326076 and T5 of BE486748. The approximate position of stop codons are indicated by asterisks. T1 and T2 may represent alternative transcripts, with T1 leading to a truncated gene product. T3 contains 28 bp that are not found in the genomic sequence and therefore most likely are artefacts. T4 and T5 probably represent unprocessed transcripts.

Figure 4 Bovine genomic sequence containing *DGAT* and parts of *HSF1* (3'end). Start codon (position 3605), stop codon (position 11906) and polyA signal (position 12163) of *DGAT* and stop codon (position 13731) and putative polyA signal (position 13439) of *HSF1* are in bold.

Figure 5 Variable PCR amplification by a, individual animals and b, pooled samples.

Figure 6 Consed views of sequencing traces for positions 10430-10437 within *DGAT* demonstrating the effect of DMSO in the PCR at variable positions 14433 and 14434 of a heterozygous animal (GC/ AA). a, three repetitions without DMSO. b, three repetitions with 5% DMSO. Average normalized amplitude values (\pm standard deviation) in a: A 1.06 ± 0.25 , A 0.61 ± 0.16 , G 0.56 ± 0.31 , C 0.21 ± 0.14 ; in b: A 0.42 ± 0.02 , A 0.22 ± 0.02 , G 1.38 ± 0.02 C 0.59 ± 0.03 .

Figure 7 Consed views of sequencing traces for positions 10430-10437 within the *DGAT* coding sequence. Positions 10433 and 10434 are variable. (a), (b) represent homozygous animals (GC/GC, AA/ AA), respectively) and (c) a heterozygous animal (AA/GC). (d) and (e) show the frequency shift between the pools FVpool12+ (breeding value milk fat % (BVMF) = $+0.729 \pm 0.045$) and FVpool12- (BVMF = -0.445 ± 0.042), (f) and (g) between pools FVpool32+ (BVMF = $+0.669 \pm 0.063$) and FVpool32- (BVMF = -0.381 ± 0.059), (h) and (i) between pools BVpool20+ (BVMV = $+0.421 \pm 0.113$) and BVpool20- (BVMF = -0.305 ± 0.057).

Figure 8 Allelic frequencies in pooled samples from animals with high (FV12+, FV32+, BV20+) and low (FV12-, FV32-, BV20-) breeding values for milk fat content at variable positions in and around *DGAT*. The numbers below the x-axis refer to the following positions (according to the numbering in Figure 3): 1, 3343; 2, 8567; 3, 8607; 4, 9284; 5, 10433; 6, 10434; 7, 11030; 8, 11048; 9, 11993; 10, 130309. The variable positions 5 and 6 are responsible for the K232A substitution, with the frequency of the A-encoding allele being indicated.

Figure 9 Alignment of the *DGAT* amino acid sequences of *Arabidopsis thaliana* (Ath), *Brassica napus* (Bna), *Perilla frutescens* (Pfr), *Caenorhabditis elegans* (Cel), *Mus musculus* (Mmu), *Rattus norvegicus* (Rno), *Ceropithecus aethiops* (Cea), *Homo sapiens* (Hsa) and two alleles of *Bos taurus* (Bta_1, Bta_2) using PILEUP of the GCG package. Sequences are assembled using BOXSHADE (http://www.isrec.isb-sib.ch:8080/software/BOX_form.html). Numbers on the left indicate amino acid positions. Red letters indicate identical amino acids. Blue letters

indicate conserved amino acids. The red arrows indicate identical lysine residues that might play a role in Acyl CoA binding. The blue arrow indicates conserved amino acids in animal species and in the bovine allele associated with high milk fat content. The lysine to alanine mutation at this position is not conservative. The alanine residue of the allele associated with low milk fat content could have a negative effect on the Acyl CoA binding capacity of DGAT.

Figure 10 Hydrophobicity plot of DGAT as assessed by Kyte-Doolittle analysis ([http:// bioinformatics.weizmann.ac.il/hydroph/plot_hydroph.html](http://bioinformatics.weizmann.ac.il/hydroph/plot_hydroph.html)). Hydrophobic regions are above the horizontal line. a Translated transcript T0 (The effect of the K232A substitution is indicated in red (K, blue; A, red)). b Translated transcript T2 (missing amino acids 230 to 251 of transcript T0).

Figure 11 Detection of the allelic variation at the nucleotide positions 10433 and 10434 of the *DGAT* gene by *CfrI*-cleavage in a 411 bp PCR product from bovine genomic DNA (primers 1532 and 1636). Cleavage by *CfrI* is diagnostic for the alanine bearing allele. Panel A, 5% DMSO in PCR reaction; panel B, PCR without DMSO. Panel A, lane 1, lane 6: homozygous for lysine variant; Panel A, lane 2, 4, 5, 7, 8, 9: heterozygous; Panel A, lane 3, 10, 11, 12: homozygous for alanine variant. Panel B, lanes 1 - 11 represent the same animals as lanes 1 - 11 in panel A. Preferential amplification of the lysine variant (nucleotides AA) over the alanine variant (nucleotides GC) prevents the detection of the alanine variant in the heterozygotes.

Figure 12 Haplotypes of *DGAT1* based on nucleotide positions 3343, 10433, 10434, 11030, 11048, 11993 determined by direct sequencing (A) and preliminary frequency estimates for the lysine (*dark*) and alanine (*light*) encoding alleles determined by RFLP assay (B). Anatolian Black is a breed indigenous of a region known as the site of domestication of the European *Bos taurus* [Medjugorac, 1994].

Figure 13 (A) Distributions of breeding values for milk fat content of Holstein-Friesian (HF), Fleckvieh (FV) and Braunvieh (BV) artificial insemination (AI) bulls born in 1990 or later. Colored areas indicate the range of the breeding values, from

which bulls were chosen for the extreme positive (+, *dark*) and negative (-, *light*) pools for HF (32 per pool), FV (32 per pool) and BV (20 per pool), respectively. HF bulls were selected among 2857 AI bulls. The mean breeding value for milk fat content of the unselected bulls was -0.148, the standard deviation was 0.284. Bulls with breeding values above 0.48 and below -0.68 were selected. The mean breeding values (\pm standard deviations) of pooled groups were as follows: HF32+, 0.622 ± 0.125 ; HF32-, -0.771 ± 0.063 . FV bulls were selected among 4070 AI bulls. The mean breeding value for milk fat content of the unselected bulls was 0.089, the standard deviation was 0.217. Bulls with breeding values above 0.5 and below -0.3 were selected. The mean breeding values (\pm standard deviations) of pooled groups were as follows: FV32+, 0.683 ± 0.153 ; FV32-, -0.454 ± 0.061 . BV bulls were selected among 656 AI bulls. The mean breeding value for milk fat content of unselected bulls was 0.006, standard deviation 0.185. Bulls with breeding values above 0.2 and below -0.2 were selected. Mean breeding values (\pm standard deviations) of pooled groups were as follows: BV20+, 0.424 ± 0.156 ; BV20-, -0.317 ± 0.096 . (B, E) Consed views of sequencing traces for positions 10430-10437 within the *DGAT1* coding sequence for individual animals (E) and DNA pools (B). (C) Allele frequency shifts. Position of variant and bases are indicated below horizontal axis. Frequencies at position 10433 are determined by genotyping individual animals by sequencing or RFLP assay. Frequencies at position 11030 and 11048 in FV + pool are determined by sequencing. The other frequencies represent estimates from sequence traces (as described in methods). Variable positions 10433 and 10434 are responsible for the K232A substitution. (D) Bars represent the frequencies of alleles with 3, 4 5, 6 and 7 repeat units in 5'-region of *DGAT1* in + pool (*dark*) and - pool (*light*) for each breed.

Figure 14 (A) Across family test statistic curve for QTL analyses of milk fat content on chromosome 14 for a Fleckvieh granddaughter design. F ratios testing for the presence of a segregating QTL are plotted for given positions along the chromosome. The marker map with distances in cM between markers is shown on the x-axis. Empirical chromosome-wide and genome-wide 1% significance levels achieved via 10,000 permutations are indicated as horizontal lines. (B) The bars

show transformed significance levels ($\log(1/p)$) of the test statistic for a segregating QTL present within each family (x-axis). The horizontal line indicates the transformed 1% significance level for a single family after correcting for multiple testing of 20 families. QTL-effects for milk fat content and their respective standard errors are shown on top of the bars for significantly segregating sires. (C) Detection of allelic variation at nucleotide positions 10433 and 10434 (K232A) of the *DGAT1* gene by *CfrI*-cleavage in a 411 bp PCR product from bovine genomic DNA of sire 1 to 16. Cleavage by *CfrI* is diagnostic for the allele encoding alanine (GC). No DNA samples were available for sires 17 to 20.

Figure 15 Haplotypes of two segregating (*Qq*) bulls. HF: Holstein-Friesian, FV: Fleckvieh. The arrows indicate the homozygous sites, implicating these variants are not causal.

Figure 16 Distribution of breeding values of sons of non segregating sires according to whether or not they have received the lysine alleles from their dams.

The examples illustrate the invention:

Example 1: Preparation of the primers

All primers used in the following procedures were designed using the Primer3 program (www.genome.wi.mit.edu/cgi-bin/primer/primer3_www.cgi). Unless indicated directly in the text, primer sequences are listed in Table 1 and Table 2.

Example 2: Radiation hybrid panel mapping

25 ng of genomic DNA from the human-hamster radiation hybrid panel Genbrige 4 (HGMP Resource Center) were amplified with one set of primers specific for the human DGAT gene (forward (1534), 5'-GAGGCCTCTCTGCCCTATG-3'; reverse (1538), 5'-TTTATTGACACCCTCGGACC-3'). PCR was performed on 84 clones of the RH-panel and analyzed by gel electrophoresis (2% agarose). PCR conditions were as follows: 10 μ l total volume containing 0.5 μ M of each Primer, 200 μ M of each dNTP, 1 μ l 10xPCR reaction puffer, 1.5 mM MgCl₂ and 0.5 U AmpliTaq polymerase (PE Biosystems). The reactions were amplified in a TGradient Thermocycler (Biometra) under following conditions: 1 cycle at 94°C for 3 min, followed by 30 cycles at 95°C for 30 sec, 60°C for 1 min, 72°C for 1 min, followed by 1 cycle at 72°C for 10 min. Positive and negative PCR assays were reported as 1 and 0, respectively, unclear assays as 2. The data were analyzed with a program provided from The Sanger Center (www.sanger.ac.uk/Software/RHserver/RHserver.shtml).

Example 3: Screening of bovine BAC library

Screening was performed by hybridization of high-density filters. A specific PCR product of 565 bp (forward primer (1599), 5'-CGAGTACCTGGTGAGCATCC-3'; reverse primer (1601), 5'-TGTGCACAGCACTTTATTGAC-3') was used as a probe for radioactive screening of the bovine RPCI-41 genomic BAC library (Warren *et al.*, 2000). PCR conditions were as follows: 20 μ l total volume containing 0.5 μ M of each Primer, 200 μ M of each dNTP, 2 μ l 10xPCR reaction puffer, 1.5 mM MgCl₂ and 1.0 U AmpliTaq polymerase (PE Biosystems). Temperature cycling were as

follows: 1 cycle at 94°C for 3 min, followed by 30 cycles at 95°C for 30 sec, 60°C for 1 min, 72°C for 1 min, followed by 1 cycle at 72°C for 10 min. Probes were labeled with 50 μ Ci of alpha[³²P]dATP using the Megaprime DNA labeling system following the manufacturer protocol (Amersham). Labeled probe was added to the filter in Church buffer and hybridized at 67°C overnight. Filters were washed twice in 2x SSC and once in 0.5x SSC + 1% SDS for 20 minutes at 63°C, respectively. Filters were exposed to Fuji NewRX film at -80°C for 5 h. Positive clones were confirmed by PCR amplification (same primer and conditions as above) and DNA sequencing.

Example 4: Sequencing from BAC-DNA

BAC-DNA was isolated using the QIAGEN Large-Construct Kit (Qiagen) following the manufacturer protocol. In the first step, primers (Table 1) for genomic walking were derived from the known bovine sequence of exon 2 (forward, 1602) and exon 3 (reverse, 1634). In addition to that, a primer (forward, 1632) was derived from the human sequence of exon 1 showing high homology to *Cercopithecus aethiops* (accession#: AF236018), *Mus musculus* (accession#: NM_010046), *Rattus norvegicus* (accession#: AF296131). Further primers were derived from the obtained sequences. Conditions of sequencing reaction were as follows: 150 ng BAC-DNA, 0.4 mM primer and 10 μ l BigDye Ready Reaction Mix (PE Biosystems) were combined in a total volume of 25 μ l. Temperature cycling were as follows: 1 cycle at 96°C for 5 min, followed by 80 cycles at 96°C for 20 sec, 55°C for 10 sec, 60°C for 4 min. DNA was precipitated with 60% isopropanol, washed with 75% isopropanol, loaded on a 36 cm WTR acrylamid gel (5.5%) on an ABI Prism 377 DNA sequencer. Sequence data were analyzed using the Phred/Phrap/Polyphred/Consed software suite (Nickerson *et al.*, 1997; Ewing and Green, 1998; Ewing *et al.*, 1998; Gordon *et al.*, 1998).

Example 5: Preparing of genomic DNA samples

DNA was prepared from bull semen. After washing with TE buffer (10 mM TrisHCl, 1 mM EDTA), cells were lysed by adding 500 μ l PK buffer (20 mM TrisHCl, 4 mM EDTA, 10 mM NaCl), 100 μ l SDS (10%), 25 μ l DTT (1 M), 60 μ l proteinase K (20

mg/ml) and incubated at 50°C overnight. Phenol/chloroform extraction was carried out in 9.5 ml VACUTAINER® tubes (#366510, Becton Dickinson). In the first step 800 µl of phenol/chloroform/isoamylalcohol (25:24:1) was added, mixed thoroughly and centrifuged for 15 min at 2000 g at RT. Traces of phenol were removed by centrifugation after adding 800 µl of chloroform/isoamylalcohol (24:1). DNA was precipitated with ethanol and resuspended in TE buffer. DNA concentration was measured using a fluorometer and adjusted to a concentration of 25 ng/µl. Quality and quantity of DNA was independently assessed through agarose gel electrophoresis and by performing PCR (primer and conditions as in Screening of bovine BAC library). Only DNA samples showing perfect results in both gel electrophoresis and PCR were used for DNA samples for individual animals and for composing pooled DNA samples.

Example 6: Comparative sequencing

Screening for variations was performed using the DNA samples of the individual animals and the pooled DNA samples in combination with several primer sets (Table 2). Each DNA sample (50 ng) was amplified in 20 µl reactions containing 0.5 µM of each Primer, 200 µM of each dNTP, 1 µl 10xPCR reaction puffer (containing 15 mM MgCl₂), 0.5 U HotStar polymerase (Qiagen). Temperature cycles were as follows: 1 cycle at 95°C for 15 min, followed by 35 cycles at 94°C for 1 min, 60°C for 1 min, 72°C for 1 min, followed by 1 cycle at 72°C for 10 min. The PCR amplified fragments were directly purified with the QIAquick PCR purification kit (Qiagen) and analyzed on a 1.5% agarose gel. Conditions of sequencing reaction were as follows: In a total volume of 10 µl was combined 20 ng PCR fragment, 0.5 µM Primer, 4 µl BigDye Ready Reaction Mix (PE Biosystems). Temperature cycling were as follows: 1 cycle at 96°C for 15 sec, followed by 25 cycles at 96°C for 10 sec, 51°C for 5 sec, 60°C for 4 min. DNA was precipitated in 60% isopropanol, washed with 75% isopropanol and run on a 36 cm WTR 5.5% acrylamid gel on an ABI Prism 377 DNA sequencer. Sequence data were analyzed using the Phred/Phrap/Polyphred/Consed software suite (Nickerson *et al.*, 1997; Ewing and Green, 1998; Ewing *et al.*, 1998; Gordon *et al.*, 1998).

Example 7: Estimation of allelic frequencies based on sequencing traces

The amplitude values at the variable positions were extracted from data files ".poly" created by the base calling program phred. The amplitude value for a given base was divided by the normalization factor for that base. The normalized amplitude value of pooled DNA (P) was compared with the amplitude value of homozygous (Ho) or heterozygous (He) individual animals or monomorphic pools. Averages were taken when amplitude values were available for more than one animal. Frequency estimates (F) were obtained by the following calculations: $F = P / Ho$ or $F = (0.5 \times P) / He$.

Example 8: RFLP-Analysis of PCR-Fragments

The genotyp of an individual or group of animals was tested by the use of RFLP-analysis. Detection of allelic variation at the nucleotide positions 10433 and 10434 of the *DGAT* gene was effected by *CfrI*-cleavage in a 411 bp PCR product from bovine genomic DNA (primers 1532 and 1636). Cleavage by *CfrI* is diagnostic for the alanine bearing allele. The result of a test is shown in figure 11. PCR reactions were carried out in the presence (panel A) or absence (panel B) of 5 % DMSO. PCR-products were isolated following common protocols as known by a person skilled in the art and incubated with the restriction endonuclease *CfrI* under conditions in line with manufactures advice. Figure 11 shows in panel A, lane 1 and lane 6 samples, which are homozygous for lysine variant. In lane 2, 4, 5, 7, 8, 9 of panel A samples with heterozygous genotype are shown. Furthermore, lane 3, 10, 11, 12: show samples which are homozygous for alanine variant. In panel B, lanes 1 - 11 samples of the same animals as shown in lanes 1 - 11 of panel A are displayed. Preferential amplification of the lysine variant (nucleotides AA) over the alanine variant (nucleotides GC) prevents the detection of the alanine variant in the heterozygotes.

Example 9: Direct sequencing reveals at least 8 haplotypes of DGAT1

Direct sequencing in animals belonging to different breeds of *Bos taurus taurus* and *Bos taurus indicus* as well as in animals of *Bos grunniens* (yak) and *Bubalus*

bubalus (water buffalo) at 6 of the variable nucleotide positions allowed to derive at least 8 haplotypes (Fig. 12). Lysine encoding haplotypes are present in yak and water buffalo. Thus, the lysine encoding variant is likely to represent the ancestral state of *DGAT1*. However, the K232A substitution is likely to have taken place early in the history of domesticated cattle or even before domestication as surmised by the presence of the alanine variant in the "old" cattle breed Anatolian Black. An RFLP assay was applied to obtain preliminary estimates on the frequency of the lysine and alanine encoding alleles in several cattle breeds and species of *Bovinae* subfamily (Fig. 12).

Example 10: Distribution of breeding values for milk fat content

The frequencies at 6 variable positions in the pools of animals with high and low breeding values for milk fat content, respectively, are visualized in Fig. 13. There are distinct differences for the Fleckvieh and Holstein-Friesian-Friesian breeds in the frequencies between the groups of animals with low and high breeding values for milk fat content, respectively, indicating association between variation in the *DGAT1* gene and genetic variation of the milk fat content. The most extreme differences are between the "low" and "high" pools in the Holstein-Friesian breed. In both breeds, the lysine encoding variant is more frequent in animals with high breeding values for milk fat content. The lysine encoding allele is also slightly less more frequent in the Braunvieh animals from the high end of the distribution of the milk fat content breeding values.

Example 11: Across family test statistic curve for QTL analyses of milk fat content on chromosome 14 for a Fleckvieh granddaughter design

Another argument for *DGAT1* (or linked loci) being responsible for the QTL-variation on chromosome 14 is provided by the results obtained from interval QTL mapping in the Fleckvieh breed using a half-sib design, the so called granddaughter design. The test statistic for the presence of a QTL along chromosome 14 (Fig. 14) indicates the

most likely position of the QTL close to marker *ILSTS039*. Evidence was highly significant for segregation of the QTL in two out of 20 families (Fig. 14). Estimates of QTL effects for milk fat content in the segregating families were found to be 0.313 ± 0.070 and 0.409 ± 0.064 , respectively. These effects greatly exceed the genetic standard deviation of 0.2 in the Fleckvieh population. The genotypes at the predicted K232A substitution determined by an RFLP assay are compatible with the heterozygous status of the segregating (*Qq*) sires and homozygosity of the alanine encoding variant of the non-segregating (most likely *qq*) sires (Fig. 14).

Example 12: Haplotypes of two segregating (*Qq*) bulls

Direct sequencing of *DGAT1* from DNA and determining the repeat number of the 5'-VNTR in the two segregating bulls and some of their progeny allowed to derive the haplotypes based on the genotypes of the homozygous progeny. The lysine encoding variant is present on two different haplotypes, i.e. the only lysine bearing haplotype in Holstein-Friesian and a Fleckvieh-specific haplotype (Fig. 12, Fig. 15). This could indicate that a lysine encoding allele has been introduced into Fleckvieh from Holstein-Friesian. Pedigree analysis indeed shows that the great-grandfather of bull 899 was a purebred Holstein-Friesian sire while there is no indication of Holstein-Friesian ancestry for bull 705. Three of the 7 variable positions that make up the haplotypes are homozygous in *Qq* bull 705 (Fig. 15). Thus they can be excluded to be causal. The variants responsible for the K232A polymorphism, however, are heterozygous in both *Qq* bulls.

Example 13: Distribution of breeding values of sons of non segregating sires

An independent association study was carried out based on the breeding values for milk fat content of the sons of non segregating sires. These sons were grouped according to the allelic variant (lysine or alanine) which they have received from

their dams as determined by the RFLP assay. The respective means of breeding values were compared after correction of half the sire's breeding value (Fig. 16). The difference of +0.265 for the group carrying the lysine variant was highly significant ($P < 0.0001$) and strongly supports the size of the gene substitution effect found via linkage analysis. It is also in agreement with the results of the association study presented above. Since the dams can be considered to represent a random sample of the Fleckvieh population with regard to milk fat content, the association involving the sons of non segregating sires is not likely to be confounded by admixture.

Example 14: Mast Experiment "Dummersdorf" – Evaluation of DGAT

Objective: Impact of DGAT for intramuscular fat content.

Material:

The experiment is based on data obtained from 56 slaughtered fattened animals of both gender of the races Deutsche Holstein Friesian ($n=29$) and Charolais ($n=27$). IMF-values of MLD (IMF_MLD) and Bratenstück [bitte übersetzen] (IMF_SEMI) and the exchange of K232A in DGAT were determined. The allelic frequency of the lysine variant, in both tested samples, were estimated as 11% for Charolais and 45% for Holstein Friesian.

Statistical analysis:

The statistical analysis was established by using the method of least squares which is part of the program SAS (Version 8.02). The analysis of the total material was based on the model:

$$Y_{ijklm} = \text{Race}_i + \text{Father}_j (\text{Race}_i) + \text{Gender}_k + \text{DGAT-Genotype}_l + e_{ijklm}$$

In another analysis, the data was evaluated for each race separately, wherein the effect of the race of the above-indicated model is left out. By employing the variance analysis, the contribution of the individual factors for the establishment of the IMF

properties was tested. Moreover, least square means were calculated for the specific genotypes, the differences of which represent an estimate reflecting the differences between these genotypes.

Results:

All experiments showed a significant gender-impact. Table 13 summarizes the F- and p-values and levels of significance (n.s: not significant; *: $p < 0.05$) of the variance analysis for the effect of DGAT genotypes. The results indicate a significant impact of DGAT on IMF_SEMI and no indication of an impact on IMF_MLD. The increased F-values of Holstein Frisian in comparison with Charolais (when data was evaluated for each race separately) may rest on the fact that a homozygous lysine variant never occurred in Charolais. From analyses on the TG locus a recessive inheritance is suggested, wherein Alanin is dominant over Lysine, thus, preventing the detection of the effect on IMF in Charolais.

Table 14 summarizes the least square means and their standard error. The predominance of L/L genotypes over L/A and A/A, as evident from the analysis, amounted to 1.6% percent in IMF_SEMI. When analyzed separately, on average a similar difference is found in Holstein Frisian. However in the latter case, the results for the genotypes L/A and A/A are less uniform and have to be discussed with caution since they are associated with a high standard error. The differences observed are of a magnitude which are likely to be only possible in extremely fastened animals. The resulting high variability of starting material may also be the reason for a lack of statistical support of the large differences in IMF_MLD of Holstein Frisian.

Tables:

Table 1: Primers used for sequencing of BAC-DNA

Location in <i>DGAT</i>	#	Direction	Sequence
5'end	1738	reverse	5'-TGATGCCTACCTAAGCTCTACC-3'
5'end	1739	reverse	5'-TTTAGGGTCTGAGCCACCAG-3'
5'end	1728	reverse	5'-TCCCGACTCTTTGTGACTCC-3'
5'end	1734	reverse	5'-TGGATTGCAAAGTCCTGTCC-3'
5'end	1717	reverse	5'-CAGGAAGGGCCTCTGTACC-3'

5'end	1716	reverse	5'-ACAGCTGGAGTGAGGACACC-3'
5'end	1710	reverse	5'-CCCTCAGCGCTAGGACTC-3'
5'end	1709	reverse	5'-TGTCTTGGAGTAGCGTGTGG-3'
5'end	1706	reverse	5'-AGGCCCCCACAGTAGACAAG-3'
5'end	1705	reverse	5'-ACGGTCGTGCTCTGTGAAC-3'
5'end	1699	reverse	5'-CCCTTGTCCTCGCTCTATAAAC-3'
5'end	1698	reverse	5'-CGCGCATACCTTTGTAGTCC-3'
5'end0	1697	reverse	5'-CGCCTCTACTACGCCACTG-3'
Exon 1	1632	forward	5'-GCCACTGGGAGCTGAGG-3'
Intron 1	1681	reverse	5'-ACAGCTGTGCACCAAGGTC-3'
Intron 1	1680	forward	5'-TGGCTGCTCTAGGGTCAAAG-3'
Intron 1	1693	forward	5'-ATCTTCACTGGGTGCTGTGG-3'
Intron 1	1694	forward	5'-CTGCTCCTGTCCTGTTGATG
Intron 1	1696	reverse	5'-AGCCACCTCATGCTACAACC-3'
Intron 1	1695	reverse	5'-GCCCTCTTCTTCATGACTCTG-3'
Intron 1	1679	reverse	5'-GGCCACCATTCAAACCAC-3'
Exon 2	1602	forward	5'-GAATTGGTGTGTGGTGATGC-3'
Intron 2	1675	reverse	5'-GGTAGGGTCCCAGGGTACG-3'
Intron 2	1673	forward	5'-GCCCACTCTGCAGGACTC-3'
Intron 2	1674	reverse	5'-CAGTCCTGCTCCCTCCAG-3'
Intron 2	1671	reverse	5'-TGACAGGCTCAGAGATGCAG-3'
Intron 2	1660	reverse	5'-AGCCCCAGTGAAGTCCAAG-3'
Exon 3	1634	reverse	5'-TAGAAATAACCGTGCGTTGC-3'
Exon 4	1633	reverse	5'-ACCTGGATGGGGTCCAC-3'
3'end	1593	forward	5'-GTGGGTGTTGGACTGCTTTG-3'
3'end	1711	forward	5'-CCATGCTCTGGAAACCCTAC-3'
3'end	1729	forward	5'-TCAGCAGGTAGTTGGGTGTG-3'
3'end	1730	forward	5'-GAAACCCTGAGGCTGTGC-3'
3'end	1732	forward	5'-CCCACCTGGTCCTCTAGTGC-3'
3'end	1733	forward	5'-CCAGGAGGCTCCAGTGTG-3'
3'end	1737	forward	5'-GTTCTGAGCCCGTCAGCAG-3'
3'end	1739	forward	5'-TTTAGGGTCTGAGCCACCAG-3'

Table 2: Primers used for PCR and comparative sequencing of genomic DNA

Location in Forward primer DGAT			Reverse primer	
	#	Sequence	#	Sequence
Exon 1	1701	5'-CGCGTTGGGTGTCAGC-3'	1681	5'-ACAGCTGTGCACCAAGGTC-3'
Exon 2	1702	5'-TGGCTTCTGCAGTGGACTC-3'	1675	5'-GGTAGGGTCCCAGGGTACG-3'
Exon 3-4	1670	5'-GTGGCTGACAGCGTTATGTC-3'	1676	5'-GTTCAGGCCCCAGATCAGC-3'
Exon 4-6	1614	5'-TATGGCATCCTGGTGGAC-3'	1617	5'-AGTGATAGACTCGAGGAGAAAGG-3'
Exon 6-7	1616	5'-GGAGCTCTGACGGAGCAG-3'	1635	5'-GTTGACGTCCCGGTAGGAG-3'
Exon 7-9	1532	5'-GCACCATCCTCTTCTCAAG-3'	1636	5'-GGAAGCGCTTTCGGATG-3'
Exon 9-11	1618	5'-CCCTGTGCTACGAGCTCAAC-3'	1678	5'-CACAGCTGGCTCCCTCAG-3'
Exon 11-14	1638	5'-GCCATCCAGAACTCCATGA-3'	1640	5'-CAGGGATGTTCCAGTTCTGC-3'
Exon 13-16	1677	5'-GAGTTCTACCGGGACTGGTG-3'	1641	5'-ATCATGCCCGGTGAAGGC-3'
Exon 16-17	1599	5'-CGAGTACCTGGTGAGCATCC-3'	1601	5'-TGTGCACAGCACTTTATTGAC-3'
5'end	1755	5'-AGAAATGGGAAGTGCAGACC-3'	1738	5'-TGATGCCTACCTAAGCTCTACC-3'
5'end	1754	5'-CAGGGTGGGATCACCTGAG-3'	1734	5'-TGGATTGCAAAGTCCTGTCC-3'
5'end	1753	5'-GGTGGATGACGGGTAGAGG-3'	1716	5'-ACAGCTGGAGTGAGGACACC-3'
5'end	1721	5'-TGAGGCCCTGATCTCTCAAC-3'	1709	5'-TGTCTTGAGTAGCGTGTGG-3'
5'end	1722	5'-AAGGGGATACTCCTGATCCAC-3'	1706	5'-AGGCCCCCACAGTAGACAAG-3'
5'end	1723	5'-TCTGCAGATGAAGGCAGAAG-3'	1698	5'-CGCGCATACCTTTGTAGTCC-3'
3'end	1711	5'-CCATGCTCTGGAAACCCTAC-3'	1718	5'-GCGGCAGAGCCAGTAGAG-3'
3'end	1729	5'-TCAGCAGGTAGTTGGGTGTG-3'	1756	5'-CTCCCTGTCTGTTCTCCTG-3'
Intron 1	1866	5'-GACACCTGGTGCGTCCTTC-3'	1867	5'-GAGGGGAGCATTTCCCAATC-3'
Intron 1	1868	5'-TACCCCCACAGACTGTCCTC-3'	1679	5'-GGCCACCATTCAAACCAC-3'
Intron 2	1602	5'-GAATTGGTGTGTGGTGATGC-3'	1674	5'-CAGTCCTGCTCCCTCCAG-3'
Intron 2	1673	5'-GCCACACTCTGCAGGACTC-3'	1671	5'-TGACAGGCTCAGAGATGCAG-3'
Intron 2	1672	5'-TGGTAAGCTGGCTGGTTAGG-3'	1634	5'-TAGAAATAACCGTGCGTTGC-3'

Table 3: Results of PCR analysis of Genebridge 4 (GB4) hamster-human radiation hybrid panel

No.	Cell line	PCR assay ^(a)	28	4G1	0
No.	Cell line	PCR assay	No.	Cell line	PCR assay
1	4A4	0	29	4G5	0
2	4A5	2	30	4G6	0
3	4AA5	1	31	4G7	0
4	4AA7	0	32	4G11	1
5	4B2	2	33	4H1	0
6	4B3	0	34	4H8	0
7	4B9	2	35	4H9	1
8	4BB1	0	36	4H12	0
9	4BB6	0	37	4I1	0
10	4BB8	1	38	4I4	1
11	4BB10	2	39	4J2	0
12	4BB12	2	40	4J5	0
13	4C3	1	41	4J9	0
14	4C11	0	42	4K5	0
15	4CC8	0	43	4K7	1
16	4D1	0	44	4K8	2
17	4D7	0	45	4K9	0
18	4DD2	0	46	4K12	1
19	4DD5	1	47	4L3	1
20	4DD8	0	48	4L4	0
21	4E2	1	49	4L6	0
22	4E4	0	50	4M4	0
23	4E6	0	51	4M5	1
24	4E11	0	52	4N3	0
25	4F6	1	53	4N5	0
26	4F7	1	54	4N6	0
27	4F13	0			

55	4N7	0
56	4N12	1
No.	Cell line	PCR assay
57	4O5	0
58	4O10	2
59	4P2	0
60	4P9	0
61	4P11	0
62	4Q2	1
63	4Q4	0
64	4R1	0
65	4R2	0
66	4R3	0
67	4R5	0
68	4R6	0
69	4R10	1
70	4R12	2
71	4S3	1
72	4S6	0
73	4S10	2
74	4S12	0

No.	Cell line	PCR assay
75	4T3	0
76	4T4	0
77	4T10	0
78	4T11	0
79	4U1	1
80	4U3	2
81	4V2	1
82	4V3	0
83	4V7	0
84	4V8	0
85	4W1	0
86	4Y4	0
87	4Y8	0
88	4Y9	0
89	4Z5	0
90	4Z6	0
91	4Z9	1
92	4Z11	0
93	4Z12	0

(a) 0, negative; 1, positive; 2, not assayed

Table 4: Bovine ESTs identified in the EST database using the human *DGAT* mRNA sequence (accession XM_005135) as input for BLASTN (Continued)

Accession	Size (in bp)	Source of mRNA	Position in bovine <i>DGAT</i> ^(a)
AW446908	479	pooled tissue from lymph node, ovary, fat, hypothalamus, and pituitary	256-780 (exon 2-9)
AW483961	205	pooled tissue from day 20 and day 40 embryos	1594-1745 (3'UTR)
AW486026	385	pooled tissue from day 20 and day 40 embryos	1336-1720 (exon17-3'UTR)
AW652329	542	pooled tissue from lymph node, ovary, fat, hypothalamus, and pituitary	990-1530 (exon 13-3'UTR)
BE664362	415	pooled tissue from day 20 and day 40 embryos	1321-1735 (exon17-3'UTR)
BE753833	422	pooled tissue from testis, thymus, semiten- dono sus muscle, longissimus muscle, pancreas, adrenal, and endometrium	1369-1745 (exon17-3'UTR)
BE664357	456	pooled tissue from day 20 and day 40 embryos	1321-1745 (exon17-3'UTR)
BE900091	527	adipose tissue	1097-1561 (exon14-3'UTR)
BE751071	475	pooled tissue from testis, thymus, semiten- dono sus muscle, longissimus muscle, pancreas, adrenal, and endometrium	1087-1560 (exon14-3'UTR)
AW446985	485	pooled tissue from lymph node, ovary, fat, hypothalamus, and pituitary	594-1143 (exon 7-11)
AW326076	141	pooled tissue from lymph node, ovary, fat, hypothalamus, and pituitary	703-772 (exon 8-9)
BE486748	174	mammary tissues at eight physiological, developmental, and disease states	906-986 (exon 11-12)

(a) Base 1 = first base of start codon

Table 5: Exon-intron structure of the bovine DGAT gene

Exon	Position in bovine <i>DGAT</i> ^(a)	Size (bp)	5'-splice donor ^(b)	Intron	Size (bp)	3'-splice acceptor ^(b)
1	1-191	191	CCTGAG g tagcg	1	3617	ctccagGTGTCA
2	192-279	88	ATGCTG g tacgt	2	1944	tcgcagATCTTA
3	280-320	41	CATCA A gtgagt	3	79	ctgcagGTATGG
4	321-406	86	TCATT G gtgagc	4	92	cctcagTGGCCA
5	407-459	53	GCCGT G gtaagc	5	215	ccccagGGAGCT
6	460-565	106	CTCCAG g tgggc	6	89	ccacagTGGGCT
7	566-679	114	AGGCT G gtgagg	7	100	tcgtagCTTTGG
8	680-754	75	ACCGC G gtgagg	8	70	ttccagATCTCT
9	755-858	104	GAGAT G gtgagg	9	90	ccccagCTGTTC
10	859-897	39	CAGCAG g tacgt	10	60 ^(c)	ttgcagTGGATG
11	898-939	42	TTCAAG g tgagc	11	73	ccacagGACATG
12	940-984	45	CTGGC G gtgagt	12	74	ccacagGTCCCC
13	985-1097	113	CTGGT G gtgggt	13	87	ccgcagGAACTC
14	1098-1163	66	CATCAG g tgggt	14	86	ccgcagACACTT
15	1164-1251	88	CACGAG g tcagt	15	81	cctcagTACCTG
16	1252-1314	63	GCGCAG g tgagc	16	72	ccccagATCCCG
17	1315-1470	156				

(a) Base 1 = first base of start codon

(b) Exon sequences are indicated in upper case letters, intron sequences in lower case letters. The consensus splice site sequences are in boldface.

(c) Intron 10 contains a (G)_n stretch that could not be resolved by sequencing.

Table 6: Panel of individual animals and animals belonging to a pool

	Lab. no.	Herdbook no.	Breed	Sub-species ^(a)
individual animals	FV19	7620	Simmental	<i>taurus</i>
	FV27	25100	Simmental	<i>taurus</i>
	FV28	50148	Simmental	<i>taurus</i>
	SB26	790580	Simmental	<i>taurus</i>
	SB37	102430	Simmental	<i>taurus</i>
	SB45	252006	Simmental	<i>taurus</i>
	AN1		Angus	<i>taurus</i>
	KE2		Kerry	<i>taurus</i>
	SA4		Sahiwal	<i>indicus</i>
	HA8		Hariana	<i>indicus</i>
SBpool	SB 2	102399	Holstein-Friesian	<i>taurus</i>
	SB 9	790121	Holstein-Friesian	<i>taurus</i>
	SB 13	790223	Holstein-Friesian	<i>taurus</i>
	SB 14	790253	Holstein-Friesian	<i>taurus</i>
	SB 22	790510	Holstein-Friesian	<i>taurus</i>
	SB 33	790361	Holstein-Friesian	<i>taurus</i>
	SB 41	790062	Holstein-Friesian	<i>taurus</i>
	SB 43	790183	Holstein-Friesian	<i>taurus</i>
	SB 44	102350	Holstein-Friesian	<i>taurus</i>
	SB 47	102315	Holstein-Friesian	<i>taurus</i>

(a) Bos taurus taurus or Bos taurus indicus

Table 7: Composition of DNA pools: Fleckvieh (Bavarian Simmental) breed

Pool ^(a)	Lab. no.	Herdbook no.	Name	Breeding value
FVpool32+	901	194100	HASTROL	0.83
	902	195260	PROLAP	0.78
	903	50223	LABTON	0.77
	906	39910	RAPID	0.75
	907	169044	HAGENT	0.74
	910	178317	LOCANDA	0.71
	911	165011	HAGER	0.70
	912	7889	ROLAND	0.70
	1066	1146	LOMBARD	0.70
	913	34380	ALPAN	0.69
	914	187217	HALLSTRAS	0.69
	916	60535	LAMBADA	0.69
	917	60250	PLANSEE	0.69
	918	54474	PROMO	0.69
	919	172162	LOMB	0.68
	920	184506	LOMO	0.68
	921	169042	HAGSON	0.67
	922	172174	LOMBOLO	0.66
	923	178308	LORETTO	0.66
	924	165010	HAGEL	0.65
	925	22153	RALBIT	0.65
	926	645073	ZEPTER	0.65
	927	60527	ALPIN	0.64
	930	34554	STREUSAND	0.63
	932	187049	HALLERTAU	0.62
	933	21784	UTNACH	0.62
	935	187138	HALBEM	0.59
	936	175061	HALLEM	0.59
	937	191053	HATARI	0.59
	939	53535	GAST	0.58
	940	191045	RODOS	0.57
	942	50246	FODA	0.56
FVpool32-	1019	45432	HONER	-0.31
	1021	53381	PRO	-0.31
	1023	178075	RAVELLI	-0.31
	1025	191283	WATL	-0.31
	1026	39733	WESPE	-0.31
	1029	68130	RAUDI	-0.33
	1032	27876	HERMANUS	-0.34
	1033	21971	HOPPE	-0.34
	1034	22043	HOPURG	-0.34
	1035	60552	HUMBACH	-0.34
	1036	68030	ZAR	-0.34

SUBSTITUTE SHEET (RULE 26)

	1038	22093	PRONER	-0.35
	1039	184256	RAUWOLF	-0.35
	1040	187114	RIVA	-0.35
	1043	184280	JUL	-0.36
	1046	53487	BONWEIN	-0.37
	1047	53493	PREUS	-0.37
	1048	68175	RAMSES	-0.37
	1049	53607	ROTWEIN	-0.37
	1050	53625	PRODOMO	-0.38
	1051	176156	RAFAEL	-0.38
	1053	27848	WIND	-0.39
	1054	68040	HIRTE	-0.41
	1055	53517	WICHT	-0.41
	1056	7787	WHISKY	-0.43
	1058	176009	FREDL	-0.45
	1060	39860	WIM	-0.46
	1061	53460	WINZER	-0.46
	1062	53293	ZECHER	-0.46
	1063	27847	RENOIR	-0.47
	1064	68195	RASTER	-0.51
FVpool12-	1065	27851	WICKY	-0.51

- (a) The bulls were selected among 4070 artificial insemination bulls born 1990 and later. The mean breeding value fat % of the unselected bulls was 0.089, the standard deviation 0.217. Bulls with breeding values greater 0.5 (N = 154, mean = 0.646 ± 0.117) and smaller -0.3 (N = 89, mean = -0.380 ± 0.062) were selected. DNA samples could be obtained from 48 bulls on the positive side (mean = 0.647 ± 0.079) and 36 bulls on the negative side (mean = -0.381 ± 0.079). The mean breeding values (\pm standard deviations) of the pooled groups were as follows: FVpool12+, 0.729 ± 0.045 ; FVpool32+, 0.669 ± 0.063 ; FVpool32-, -0.381 ± 0.059 ; FVpool12-, -0.445 ± 0.042

Table 8: Composition of DNA pools: Braunvieh (Brown Swiss) breed

Pool ^(a)	Lab. no.	Herdbook no.	Name	Breeding value
BVpool120+	909	78780	BREILORI	0.73
	929	79030	BREICON	0.63
	943	340530	EURO	0.54
	951	79195	VINCOL	0.50
	952	79115	EMOZ	0.47
	953	348544	STRIFMAN	0.46
	954	78475	DOTRAY	0.45
	955	348105	BRAY	0.44
	956	349447	BREIMORY	0.42
	957	78635	DOTION	0.40
	959	77888	ROMEIS	0.38
	961	348247	BREIZ	0.37
	962	348591	STRIZIN	0.37
	964	349569	HUCNOS	0.35
	965	72695	DOLEIN	0.34
	966	340573	BREISAD	0.33
	967	340015	STRELE	0.32
	968	78980	EMPIKT	0.31
	971	79080	RELVIN	0.31
	972	78880	BAYDOT	0.29
BVpool120-	1004	78225	DOBROY	-0.22
	1006	78200	VISTAR	-0.22
	1007	348215	CREVIN	-0.24
	1008	72625	TRALAS	-0.24
	1009	348607	VIVAT	-0.24
	1011	72680	BAGAT	-0.27
	1012	72470	SIRAY	-0.27
	1014	72930	PETOS	-0.29
	1015	78090	SIMPUR	-0.30
	1017	78470	BARI	-0.31
	1018	78840	BLESTRI	-0.31
	1024	78860	RENTZ	-0.31
	1027	78560	JETSTRI	-0.30
	1028	72490	JUP	-0.30
	1030	85550	RESTOR	-0.30
	1037	78015	DUKE	-0.40
	1042	78695	CRAUTS	-0.40
	1044	348104	PETMAN	-0.40
	1045	340010	BAY	-0.40
	1057	78155	JARGI	-0.40

(a) The bulls were selected among 656 artificial insemination bulls born 1990 and later. The mean breeding value "fat %" of the unselected bulls was 0.006, the standard deviation 0.185. Bulls with

breeding values greater 0.2 (N = 84, mean = 0.325 ± 0.108) and smaller -0.2 (N = 56, mean = -0.334 ± 0.101) were selected. DNA samples could be obtained from 54 bulls on the positive side (mean = 0.316 ± 0.111) and 22 bulls on the negative side (mean = -0.306 ± 0.055). The mean breeding values (\pm standard deviations) of the pooled groups were as follows: BVpool20+, 0.421 ± 0.113 ; BVpool20-, -0.305 ± 0.057 .

Table 9: Variable positions in and around *DGAT* and genotypes of individual animals

Position	Variation	Animals									
		FV19	FV27	FV28	SB26	SB37	SB45	AN1	KE2	SA4	HA8
1465-1554	4, 5, 6 ^(a)	4,4	4,4	4,4	5,5	5,6		5,6		4,4	5,6
3343	C - G	CC	GC	CC	CC	CC	CC	CC		CC	CC
3399	T - G	TT	TT	TT	TT	TT	TT	TT		GG	TG
7232	A - G	AA	AA	AA	GG	AA	AA	AA		GG	GG
8567	A - G				AA						
8607	G - A				GG						
9284	T - C ^(b)										
10147	A - C	AA	AA	AA	AA	AA	AA	AA		CC	AA
10433	G - A	GG	GG	GG	AA	AG	AG	GG	GG	AA	AA
10434	C - A	CC	CC	CC	AA	CA	CA	CC	CC	AA	AA
10508-10512	G5 - G6	G5G5	G5G5	G5G5	G5G5	G5G5	G5G5	G5G5	G5G5	G5G5	G5G6
ca. 10800	PCR ^(c)	-	-	-	+	+	+	+	+/-	+	+
11030	G - A	GG	GG	GG	AA	AG	AG	GG	GG	GG	AA
11048	C - T	TT	TT	TT	CC	CT	CT	TT	TT	CC	CC
11993	T - C	TT	TT	TT	CC	TC	TC	TT	TT	TT	TT
12005	A - C	AA	AA	AA	AA	AA	AA	AA	AA	CC	AA
12036	T - C	TT	TT	TT	TT	TT	TT	TT	TT	CC	TT
12056	A - G	AA	AA	AA	AA	AA	AA	AA	AA	GG	AA
12136	G - A	GG	GG	GG	GG	GG	GG	GG	GG	AA	GG
13309	G - Cb										

(a) Number of repeats (AGGCCCCGCCCTCCCCGG)

(b) Detected in pooled DNA (see Table 8)

(c) Variable PCR amplification (+, PCR product; -, no or very weak PCR product)

Table 10: Repeat at position 1465-1554 and genotypes of pooled samples

4,4 ^(a)	4,5 ^(a)	5,5 ^(a)
FV12-	FV12+	
FV32-	FV32+	
	BV20-	BV20+

(a) Number of repeats (AGGCCCGGCCCTCCCCGG)

Table 11: Allelic frequencies estimated from sequencing traces of pooled samples

Position ^(a)	Exchange	SBpool	FV12+	FV12-	FV32+	FV32-	BV20+	BV20-
3343	C – G	1	1	0.79	1	0.70	1	0.82
8567	A – G	n.d.	n.d.	n.d.	0.42	0	n.d.	n.d.
8607	G – A	n.d.	n.d.	n.d.	0	0.49	n.d.	n.d.
9284	T – C	n.d.	n.d.	n.d.	0.54b	0.92d	0.90b	1b
10433	G – A	n.d.	0.39 ^(b)	1b	0.46b	1	0.90b	1b
10434	C – A	n.d.	0.36b	1b	0.41b	1	0.93b	1b
11030	G – A	n.d.	0.68b	1b	0.64b	1	1b	1b
11048	C – T	n.d.	0.48b	0.20b	0.48b	0.26d	0b	0b
11993	T – C	0.61	0.64	1	0.65	1	1	1
130309	G – C	n.d.	n.d.	n.d.	0.39	1	1	n.d.

(a) Only positions with single base exchanges and that are variable within *Bos taurus taurus*

(b) 5% DMSO in PCR

Table 12: Genotypes of individual animals

Pool	Lab #	Breeding value	Position (base)			
			10433 (A) ^(a)	10434 (A) ^(a)	11030 (A) ^(a)	11048 (C) ^(a)
FV12p+	901	0.83	1	1	0	0
	902	0.78	0	0	-	-
	903	0.77	1	1	0	0
	906	0.75	2	2	2	2
	907	0.74	1	1	0	1
	910	0.71	1	1	0	0
	911	0.70	1	1	0	1
	912	0.70	1	1	1	1
	1066	0.70	1	1	0	2
	913	0.69	1	1	0	1
	914	0.69	0	0	0	0
	916	0.69	2	2	1	2
FV32p+	Average / Frequency		0.5%	0.5%	0.18%	0.45%
	917	0.69	2	2	0	1
	918	0.69	1	1	0	1
	919	0.68	1	1	0	1
	920	0.68	0	0	0	1
	921	0.67	0	0	0	0
	922	0.66	1	1	0	-
	923	0.66	0	0	0	1
	924	0.65	1	1	0	1
	925	0.65	1	1	1	1
	926	0.65	0	0	0	2
	927	0.64	1	1	1	1
	930	0.63	1	1	1	1
	932	0.62	2	2	0	1
	933	0.62	2	2	1	0
	935	0.59	1	1	0	1

SUBSTITUTE SHEET (RULE 26)

936	0.59	1	1	0	1
937	0.59	1	1	0	1
939	0.58	1	1	1	1
940	0.57	0	0	0	1
942	0.56	1	1	0	1
Average / Frequency		0.47%	0.47%	0.15%	0.94%

(a) 0, 1, 2, number of indicated allele; - assay failure

Table 13: F- and p- values of the variance analysis

Race	IMF_MLD		IMF_SEMI	
	F-Vaule	Sig.	F-Value	P
Total	0,19	0,827 n.s.	3,47	0,040*
Holstein-Friesian	0,36	0,704, n.s.	5,35	0,013*
Charolais	0,15	0,703, n.s.	1,13	0,301, n.s.

Table 14: Least square means and standard error

animals	IMF_MLD		IMF_SEMI	
	LSM +/- s.e.		LSM +/- s.e.	
gesamt				
L/L	5,57	0,99	3,95	0,59
L/A	5,05	0,49	2,35	0,29
A/A	4,88	0,41	2,35	0,24
Holstein-Friesian				
L/L	7,07	1,04	4,33	0,53
L/A	6,14	0,61	2,39	0,31
A/A	6,08	0,82	3,04	0,41
Charolais				
L/A	3,80	0,62	2,46	0,50
A/A	3,53	0,32	1,85	0,26

References

- Asil Memisoglu (2001) Pharmacogenomics, published on www.thebiotechclub.org/Tech/pharmacogenomics.html.
- Barendse WJ (1999). Assessing lipid metabolism. International Publication Number WO 99/23248.
- Coppieters W, Riquet J, Arranz JJ, Berzi P, Cambisano N, Grisart B, Karim L, Marcq F, Moreau L, Nezer C, Simon P, Vanmanshoven P, Wagenaar D, Georges M (1998) A QTL with major effect on milk yield and composition maps to bovine chromosome 14. *Mammalian Genome* 9: 540-4.
- Ewing B, Green P (1998) Base-calling of automated sequencer traces using phred. II. Error probabilities. *Genome Research* 8: 186-194.
- Ewing B, Hillier L, Wendl MC, Green P (1998) Base-calling of automated sequencer traces using phred. I. Accuracy assessment. *Genome Research* 8: 175-185.
- Farese RV, Jr., Cases S, Smith SJ (2000) Triglyceride synthesis: insights from the cloning of diacylglycerol acyltransferase. *Curr Opin Lipidol* 11: 229-34.
- Georges M, Nielsen D, Mackinnon M, Mishra A, Okimoto R, Pasquino AT, Sargent LS, Sorensen A, Steele MR, Zaho X, Womack JE, Hoeschele I. (1995) Mapping of quantitative trait loci controlling milk production in dairy cattle by exploiting progeny testing. *Genetics* 139: 907-920.
- Goddard ME, Wiggans GR (1999). Genetic improvement of dairy cattle. *The Genetics of Cattle*. R. Fries and A. Ruvinsky. Wallingford, CABI Publishing: 511-37.
- Gordon D, Abajian C, Green P (1998) Consed: a graphical tool for sequence finishing. *Genome Research* 8: 195-202.
- Harlow and Lane (1988), "Antibodies, A Laboratory Manual", CSH Press, Cold Spring Harbor.
- Heyen DW, Weller JI, Ron M, Band M, Beever JE, Feldmesser E, Da Y, Wiggans GR, VanRanden PM, Lewin HA (1999) A genome scan for QTL influencing milk production and health traits in dairy cattle. *Physiological Genomics* 1: 165-175.
- Higgins and Hames (eds.), "Nucleic acid hybridization, a practical approach", IRL Press, Oxford 1985
- Melani, C., Rivoltini, L., Parmiani, G., Calabretta, B., Colombo, MP. (1991) Inhibition of proliferation by c-myc antisense oligodeoxynucleotides in colon adenocarcinoma cell lines that express c-myc. *Cancer Res.* 51: 2897-2901
- Nickerson DA, Tobe VO, Taylor SL (1997) PolyPhred: automating the detection and genotyping of single nucleotide substitutions using fluorescence-based resequencing. *Nucleic Acids Res* 25: 2745-51.

Rebeiz M, Lewin HA (2000) Compass of 47,787 cattle ESTs. *Anim Biotechnol* **11**: 75-241.

Riquet J, Coppieters W, Cambisano N, Arranz JJ, Berzi P, Davis SK, Grisart B, Farnir F, Karim L, Mni M, Simon P, Taylor JF, Vanmanshoven P, Wagenaar D, Womack JE, Georges M (1999) Fine-mapping of quantitative trait loci by identity by descent in outbred populations: application to milk production in dairy cattle. *Proc Natl Acad Sci U S A* **96**: 9252-7.

Sambrook et al., (1989) "Molecular Cloning, A Laboratory Manual"; CSH Press, Cold Spring Harbor.

Scopes, (1982), Protein Purification, Springer-Verlag, N.Y.

Smith SJ, Cases S, Jensen DR, Chen HC, Sande E, Tow B, Sanan DA, Raber J, Eckel RH, Farese RV, Jr. (2000) Obesity resistance and multiple mechanisms of triglyceride synthesis in mice lacking Dgat. *Nat Genet* **25**: 87-90.

Steinecke, (1995), Ribozymes, *Methods in Cell Biology* **50**, Galbraith et al. eds Academic Press, Inc. 449-460.

Threadgill DW, Fries R, Faber LK, Vassart G, Gunawardana A, Stranzinger G, Womack JE (1990) The thyroglobulin gene is syntenic with the MYC and MOS proto-oncogenes and carbonic anhydrase II and maps to chromosome 14 in cattle. *Cytogenetics and Cell Genetics* **53**: 32-36.

Velmalu RJ, Vilkkilä HJ, Elo KT, de Koning DJ, Maki-Tanila AV (1999) A search for quantitative trait loci for milk production traits on chromosome 6 in Finnish Ayrshire cattle. *Anim Genet* **30**: 136-43.

Warren W, Smith TP, Rexroad CE, Fahrenkrug SC, Allison T, Shu CL, Catanese J, de Jong PJ (2000) Construction and characterization of a new bovine bacterial artificial chromosome library with 10 genome-equivalent coverage. *Mamm Genome* **11**: 662-3.

Zhang Q, Boichard D, Hoeschele I, Ernst C, Eggen A, Murkve B, Pfister-Genskow M, Witte LA, Grignola FE, Uimari P, Thaller G, Bishop MD (1998) Mapping quantitative trait loci for milk production and health of dairy cattle in a large outbred pedigree. *Genetics* **149**: 1959-73.

CLAIMS

1. A nucleic acid molecule encoding a bovine acyl CoA:diacylglycerol transferase (*DGAT*) contributing to or indicative for low fat content of milk and to low meat marbling (intramuscular fat content) wherein said nucleic acid molecule is selected from the group consisting of:
 - (a) a nucleic acid molecule having or comprising the nucleic acid sequence of SEQ ID NO: 1;
 - (b) a nucleic acid molecule comprising the coding sequence of the polypeptide of SEQ ID NO: 2;
 - (c) a nucleic acid molecule the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (a) or (b), wherein said nucleic acid molecule has at the position corresponding to position 10433 and 10434 of the *DGAT* gene (SEQ ID NO: 1) a guanine and a cytosine residue; and
 - (d) a nucleic acid molecule the complementary strand of which hybridizes under stringent conditions to the nucleic acid molecule of (a) or (b), wherein said nucleic acid molecule has at the *DGAT* gene (SEQ ID NO: 1) position
 - (i) 3343 a cytosine, 10433 a guanine, 10434 a cytosine, 11030 a guanine, 11048 a cytosine and 11093 a thymine;
 - (ii) 3343 a cytosine, 10433 a guanine, 10434 a cytosine, 11030 a guanine, 11048 a thymine, and 11093 a thymine; or
 - (iii) 3343 a guanine, 10433 a guanine, 10434 a cytosine, 11030 a guanine, 11048 a thymine and 11093 a thymine.
2. The nucleic acid molecule of claim 1 which is mRNA, genomic DNA (gDNA) or cDNA which is derived from said mRNA by reverse transcription of said mRNA.
3. The nucleic acid of claim 2, wherein said gDNA is a gene.

4. A fragment of the nucleic acid molecule of any of claims 1 to 3 having at least 14 nucleotides wherein said fragment comprises nucleotide position 10433 and 10434 of SEQ ID NO: 1.
5. A nucleic acid molecule which is complementary to the nucleic acid of any of claims 1 to 4.
6. A vector comprising the nucleic acid molecule of any one of claims 1 to 5.
7. The vector of claim 6 comprising regulatory elements for expression of said nucleic acid molecule.
8. A primer or primer pair, wherein the primer or primer pair hybridize under stringent conditions to the nucleic acid molecule of any of claims 1 to 5 comprising nucleotide position 10433 and 10434 of SEQ ID NO: 1 or the complement strand thereof.
9. A host cell which contains the expression vector of claim 7.
10. A method for production of a functional bovine *DGAT* or a functional fragment thereof comprising:
 - (a) culturing the host cell of claim 9 containing the expression vector which comprises the nucleic acid molecule of any of claims 1 to 3 under conditions allowing the expression of the encoded polypeptide; and
 - (b) collecting the polypeptide from the culture.
11. A functional bovine *DGAT* polypeptide or a functional fragment thereof encoded by a nucleic acid molecule according to any of claims 1 to 3 or produced by the method of claim 10.

12. An antibody which binds to an epitope of the polypeptide or fragment of SEQ ID NO: 2 the epitope comprising a alanine at position 232 but not to a polypeptide or a fragment of SEQ ID NO: 4 having a lysine at position 232.
13. An antibody which binds to an epitope of the polypeptide or fragment of SEQ ID NO: 4 the epitope comprising a lysine at position 232 but not to a polypeptide or a fragment of SEQ ID NO: 2 having a alanine at position 232.
14. A transgenic, non-human animal comprising at least the nucleic acid molecule of any of claims 1 to 3 or 5.
15. The transgenic, non-human animal of claim 14 wherein said animal belongs to cattle.
16. A method of testing a mammal for its predisposition for fat content of milk and/or its predisposition for meat marbling comprising analyzing the nucleic acid of a sample comprising the gene encoding *DGAT* or a corresponding mRNA for nucleotide polymorphisms which are connected with said predisposition.
17. The method of claim 16, wherein the nucleic acid molecule analyzed is the nucleic acid molecule of claim 1.
18. The method of claim 16 wherein said nucleic acid is DNA.
19. The method of claim 18 wherein said DNA is gDNA.
20. The method of claim 16 wherein said nucleic acid is cDNA which is derived from said mRNA by reverse transcription of said mRNA.
21. The method of any of claims 16 to 20 wherein the nucleotide polymorphisms are located in the coding region of the *DGAT* gene.

22. The method of claim 21 wherein the nucleotide polymorphisms in the coding region of the gene encoding *DGAT* result in substitution, deletion and/or addition of at least one amino acid in the amino acid sequence of the polypeptide which is encoded by said gene.
23. The method of any of claims 16 to 22 wherein said nucleic acid molecule has at the position corresponding to position 10433 and 10434 of the *DGAT* gene (SEQ ID NO: 1) a guanine and a cytosine residue which correlate with a predisposition for low fat content of milk and to low meat marbling.
24. The method of claim 23, wherein said nucleic acid molecule has at the position corresponding to position:
- (a) 3343 a cytosine, 10433 a guanine, 10434 a cytosine, 11030 a guanine, 11048 a cytosine and 11093 a thymine;
 - (b) 3343 a cytosine, 10433 a guanine, 10434 a cytosine, 11030 a guanine, 11048 a thymine, and 11093 a thymine; or
 - (c) 3343 a guanine, 10433 a guanine, 10434 a cytosine, 11030 a guanine, 11048 a thymine and 11093 a thymine
- which correlates with a predisposition for low fat content of milk and low meat marbling.
25. The method of claims 16 to 22 wherein said nucleic acid molecule has at the position corresponding to position 10433 and 10434 of the *DGAT* gene (SEQ ID NO: 3) two adenine residues which correlate with a predisposition for high fat content of milk and high meat marbling.
26. The method of claim 25, wherein said nucleic acid molecule has at the position corresponding to position:
- (a) 3343 a cytosine, 10433 an adenosine, 10434 an adenosine, 11030 an adenosine, 11048 a cytosine and 11093 a thymine;
 - (b) 3343 a cytosine, 10433 an adenosine, 10434 an adenosine, 11030 an adenosine, 11048 a cytosine and 11093 a cytosine;

- (c) 3343 a cytosine, 10433 an adenosine, 10434 an adenosine, 11030 a guanine, 11048 a cytosine and 11093 a thymine;
 - (d) 3343 a thymine, 10433 an adenosine, 10434 an adenosine, 11030 a guanine, 11048 a cytosine and 11093 a cytosine;
 - (e) 3343 a cytosine, 10433 an adenosine, 10434 an adenosine, 11030 a guanine, 11048 a cytosine, and 11093 a cytosine; or
 - (f) 3343 a thymine, 10433 an adenosine, 10434 an adenosine, 11030 a guanine, 11048 a cytosine, and 11093 a cytosine
which correlates with a predisposition for high fat content of milk and high meat marbling.
27. The method of any of claims 16 to 20 wherein the nucleotide polymorphisms are located in a region which is responsible for the regulation of the expression of the product of the gene encoding *DGAT*.
28. The method of any of claims 16 to 27 wherein the nucleotide polymorphisms are single nucleotide polymorphisms (SNP).
29. The method of any of claims 16 to 28, wherein said testing comprises hybridizing the nucleic acid molecule of claim 5 as a probe under stringent conditions to the nucleic acid molecules comprised in said sample and detecting hybridization.
30. The method of claim 29 further comprising digesting the product of said hybridization with a restriction endonuclease and analyzing the product of said digestion.
31. The method of claim 29, wherein said probe is detectably labeled.
32. The method of any of claims 16 to 28, wherein said testing comprises determining the nucleic acid sequence of at least a portion of said nucleic acid molecule.

33. The method of claim 32, wherein the determination of the nucleic acid sequence is effected by solid-phase minisequencing.
34. The method of any of claims 16 to 28 further comprising, prior to analyzing the nucleic acid, amplification of at least a portion of said nucleic acid.
35. The method of claim 34, wherein in the amplification reaction at least one of the primers employed in said amplification reaction is the primer of claim 8 or belongs to the primer pair of claim 8, comprising assaying for an amplification product.
36. The method of claim 34 or 35 wherein said amplification is effected by or said amplification is the polymerase chain reaction (PCR).
37. The method of any of claims 16 to 28 or 34 wherein the nucleic acid is analyzed by the use of:
 - (a) a primer extension assay;
 - (b) a differential hybridization assay; and/or
 - (c) an assay which detects allele-specific enzyme cleavage.
38. A method of testing a mammal for its predisposition for fat content of milk and/or its predisposition for meat marbling comprising:
 - (a) preparation of a tissue sample from the subject;
 - (b) contacting the sample with an antibody of claim 12 or 13; and
 - (c) detecting whether a specific binding of said antibody to its antigen has occurred.
39. The method of claim 36 wherein binding of the antibody of claim 12 indicates a predisposition of the mammal for low fat content of milk and to low meat marbling.

40. The method of claim 38 wherein binding of the antibody of claim 13 indicates a predisposition of the mammal for high fat content of milk and to high meat marbling.
41. The method of any of claims 16 to 40, wherein the sample is isolated from cloven hoofed animals.
42. The method of claim 41, wherein the cloven hoofed animals are cattle, buffalos, yaks or pigs.
43. A kit comprising at least the fragment of claim 4, the nucleic acid molecule of claim 5, the primer or primer pair of claim 8, or one of the antibodies of claim 12 or 13 in one or more container.

1/21



Figure 1

2/21




Figure 2

3/21




Figure 3

4/21

1 CTGCCCCGAC AGGCCTGACA ACCAACAACA AGCCTTCCTC AATGCCACTA
 51 GAGAAATGGG AAGTGCAGAC CCCTTCTGTC AGCCTGCTTT CCACATCCTG
 101 ACTTCCAGAT TCAGGGGACA TGTCCCACA CTGAGGAGGC TTTCTTGGT
 151 AGCTGGAACA GGCTGGTTGT GGGGAGGAGA TACCAAGGA ATAAGAACCT
 201 CCCATGGCCA CCCCCAGCCC TTAGGCTCTA GACAGGGTGA GTCAAGTTGA
 251 GAAGATGAAT GGCAGGGCTG TGCTGGGCTC AGACAACCAA GGAACATAGA
 301 CTCCTGCCCC AGCAATGCGC GTTGGTAACG AGCAGCTAG GGTTCAGCTA
 351 AGAGGGTCCA AATCTTTCGA GACATGTGGT CAAACTGGAT CAGCCAGGGG
 401 CCAGCACAGC TGTCTGCACC CTGGCAGGGG ACAGGCCAC CAGACTCCAC
 451 TGGTGTGGAC AGCAGGAAAG CCTGAOCTGC AGTAGACCTG CTGCTTCAGG
 501 GTGGGATCAC CTGAGGTGGG CACCCCTTC TGGGGAGCAC TGTCAGCCTT
 551 CATAACCTCA GGATGAAAGC CCCAGTATT GGTAGAGCTT AGGTAGGCAT
 601 CATTGCCCAA TCTGCATATG AAGAGTCTGA CCTCAGGGA GAGAAGCAGC
 651 TTGCCAAGGG CTGCCTTGA CTTAAGCCCT GCTCCAGTTG GGCTTCCCTG
 701 GTGGCTCAGA CCCTAAAGAA TCTGCCTGCA ATGTGGGAAA CCTGGGTTC
 751 GTCCCTGGGA CGGGAAGATC CCCTGGAGAA GGGATGGCAA CCCACTCCAG
 801 TGTTCTTGCC TGAGAATCCC ACGGACAGAG GAGCCTGGCG GGCTGCAGTC
 851 CATGGAGTCG CAAAGAGTCG GACACGACTG AGCAACTAAC ACTTTCACCT
 901 TCTGCCCCAA TACCCACCCC ATCTGAACCT GAATACCTGA GTGGGTCCCA
 951 CTGGCAGGAA GAGAGGCTCC TAGAGGCCCA GTCCTCCCCA AGGCTCCTCA
 1001 GCTTTGGGGC CTGGATTGAC TGTTCAGGA CTCTGATGGG CGGCTGGGGT
 1051 GGATGACGGG TAGAGGCTGC CTCCCAGTG ACTGGGACAG GCCTAGCCTT
 1101 GTCTCCACAG GTGTCCATGG ACAGGACTTT GCAATCCAGA GGATGGGTGG
 1151 TGTGGTGCAG GCTGCTGACC ACTGTGTCCA GGGTCTTCTC TCACGGGCCC
 1201 AAGGCGCCTC CAACCTGGAG TCAGCCCAAG GCTCTTTCTA AATCCCCAAA
 1251 CCCTTCCAGC CCTTCATTCC GCCAGCTGC AGATTCTCTG TCCCAAGACA
 1301 GATGTTGCTT CCACCAGGGG GAGATTCTCT ATTGAGCTTT CTTCAACAA
 1351 CTCTCAAGC ACATTTGTCC CAAAAGACC CCACCTATCT TGACGTTTTC
 1401 CTTCTGCTT CTTCTGCTG ACCCTGSCAG CACCTCAATC AGGATCCAGA
 1451 GGTACCAGGG CTGTTCGCGC GAGGCTGCTG GAGGCTGCTG GAGGCTGCTG
 1501 AGGCTGCTGCT GAGGCTGCTG GAGGCTGCTG GAGGCTGCTG GAGGCTGCTG
 1551 AGGAGGCCCG CGCCCTGTAT CAACCTTGA CCCCCTCTTC CTCAAACAGG
 1601 CCCC GCCCGC CTTGGTACA GAGGCCCTTC CTGATTGGTG CTTTACAGT
 1651 CCGTGCCCTT TCATTGGCTT GAGGCCCTGA TCTCTCAACT CCAGCGGTGG
 1701 AACCCTTGGT TCCCTCACGT CCGGGTCAG ATCGGTTCTC TTTGATGACC
 1751 CTCGGCCAC CCTGGTGTCC TCACTCCAGC TGTTCATGT TAGCCGAAGG
 1801 CAAAGGAGCC TGGACGCGGA CACAGGGAGC CGCCCCAAC ACGTACCTTC
 1851 ACTCGTCAGT GGCTACTGTG CTCAGCTCT CCAGGCCAAC AGGCAGCCTG
 1901 AGCCGTCAAT CTTCTCTCT GCCAATCAGC GCGCCAGCCA GGCTGGCCCT
 1951 CTAGTCAGGG CTCGGTACTG AAGGATGGCA AGTCCGAAG GCTCCAGGG
 2001 ACGCGTGCGC ACGGGTTAGG GGGCTTCCCA CCAGCTGCCT GGGAGAGGGA
 2051 TAGGGAGGGA AAGGCAGAGC TCCCGGACT CAGCCCTGCT GCGCGTTCCT
 2101 GAGAGGACTC TCTCTCTCT CCATCCTCCC TTGGGAGCTA TACTGAGTCC
 2151 TAGCGCTGAG TGGCCCAACT CTGCCTATGA ATAGACGAAG GTGCTTGGAC
 2201 ACTGGCTAAG GGGATACTCC TGATCCACCG AGGCGGGGCC TGTGAGGAGG
 2251 CAAGAGGGGT TCTCCAGCCT GATGAGGTCG CTCGAGCCCT TCCACACGCT
 2301 ACTCCAAGAC ACGGGCCAGG TAGCTCCAGC CTSCAGGTA AGGATGTCAG
 2351 GCTGGCCTCA GCCGCAATG GTCCAGTGGG AGAATATGTC ACCAGGGTCC
 2401 CAGGTGCCTG TTGGTTGAGG TAAGAGGGTC AGGAGCGAGT CCGCAGGAA

Figure 4

5/21

2451 GGAGGCTTGA TCTCAGGCTG AGCCTCTTGG TTTATTTGCT TTCAGAGAGG
2501 CGGTCTTCCC AGCTTTGCTT ACCCCATGGG AGTGAACGGA GTGGGTCTG
2551 TGGCTAGGGG TGTTTCTTGT GTAAACCAGG CCTAAACTCC CGGTGAACCC
2601 TCGCATCTGG AGATCCAGGA TACTCACACT CCATGCTCTT TGCCAAATGT
2651 TTGTGAAACC AAGTAAGATC GGCCTTGCCC GCGCACGGGC CTCCTGTGTC
2701 AGTTGTTTTG GTGTATTGGT TGCTTCATTC AACGACTGGA TGA CTGCCCAG
2751 CTGTGCAATG AAACAGAAAC CTCTGGGTCC CTGCGAATCA ACACCCAGG
2801 ATCCTAACTC CCTGGCAAAA CTGGCCCAAG TGGGGAAGGC GGGAAAGTCT
2851 GCAAGTCTGC AGATGAAGGC AGAAGCGGGG CGGGTGGAGA GGCGGGCTGG
2901 CTTGTCTACT GTGGGGGCCT GGGCAGGGGA GAGGTGGCCA CCCTGGGAAT
2951 AGGTGGGCAT GGCACAAGTC CCGGAATGCG AGGACTGCGG CCTTTCTCCC
3001 CCTCCGTTCT CTGACCTGGC GCGTGTTTGA ACAGCCTAAG TGGAGGAAAA
3051 GTGGGTGCCT ACGGTGGTAA TTAGTGGGT CACAGAGCAC GACCGTGCCG
3101 CGGGATGTAC GTTCGGTAGA CGCGTTGGGT GTCAGCCTGA CGTTAACGCA
3151 CTAGGCATTT CATAAATAAC TACAAOCCCA AATTCTGCGC CTGAGCTGAG
3201 AAATGACGAA ATCCTGTGTT TATAGAGCGG GACAAGGGGC AGGCAGCGGT
3251 CAGCAGAGGC TTGTTTGAG CTGCCCGGAA GCCCCGCGTG TTCCTCGTCT
3301 GTCCGGGATT GCATTTGCCA GGAGACCACA ACTCCCAGGG TGACCGCGC
3351 GCCAGCGGAC TACAAAGTA TGC GCGCCGC GGCCCTGGGC CAGTTAGCTG
3401 TCCCGGGAAC TACGCTTCCC AGGACTCCGA GAGGAGCCGT CCGGCACGGA
3451 TTTGCACGCG CTGATTGGCG GCGCGGACCA CGGCAGTGGC GTAGTAGAGG
3501
3551
3601
3651
3701
3751 GTAGC
3801 GGTGCGCGTG ACCCCTAACC TTTGACCCCT GATAOGGGGC CCCTGCGACC
3851 CAACCTGGTG GCCCAGGCCT GTCGGCGGCA GCTCGGGCTC GAGTCCGAGA
3901 TCTGGCGCC TGGACCTTGG TGCACAGCTG TGCCCTCGG GCCTCCACGG
3951 GGAAACTTAG CGGGAGGTTG GGGGCGGAGG GTCTCCTGCC CGGAACACCC
4001 AGGTACGGGG GCCGAGGGGA GGGCAGCGGC TCAACTTCTA GACGCCCTCC
4051 CTCTGCCCTT CTTTGGTGGG TTCTGAAGCT TTCCCAGGGT GAGCCACTA
4101 CGCACAGTGT CCTCTACCTG GAAGGAGATA CAGGGGTCTT TCCTGAGGGC
4151 TATGAGGGGT GCCTTGTGGG TTGATAAAGC TCCCGGGGGA GGAGGGTGGA
4201 CCGGCGGAGA ACAGAGGCAG GGGCAGTGCG AGGGGATTTT TCATCCCTCG
4251 CAGACCCTCC AGAGAAATGGT CTTACAAAG GTCCCTCATC CGTCACCCGG
4301 CGATTGACTG GCCTAGGATC CTGCTTATTA CCAGCACAAA TGGCTGCTCT
4351 AGGGTCAAAG TGGGTCTGT AATGGGACCC TCACCCCTGG TTGGGGTACA
4401 GGGGAGGAGT TGGAAGTGCG CACACCCACA GGTGGGCGCC CTGCTTAGCT
4451 GAAGGACTGA TGGGAAGGAG TTGGGGGAGC AAGCTGCGGC TGAAAGGGAG
4501 GATCTGACCC ACGTGGGCAT CAGCTAAGTC CTGCTGGCTG CCTOCAGGCG
4551 CCCCCTTTGC CATCCTCCAC GCCCCTCCCC CCAGCCCTGA CCTTCATCCT
4601 GGTCAAGGGC TCTCAGGGGC TCTGGTTTTG GGATCAGCTC CAGAGCTAGA
4651 GGTATCAAG GAGGAAGTGG GCAACAGGTC AGTCAGCAAG GATTTGCTAT
4701 CTTCACTGGG TGCTGTGGGG AGGGGAGGGA CAAGGGCAGT TGGGGTGAG
4751 GCACTGTCCC TGCCCTTGGG GGGCACACAG TTCACTGAG AGATAAGATA
4801 GCCGCAGCCC TGAAGAGTGA GAGCAAAGGT CAGGCACAGA GTTACAGGATG
4851 ACACCAGGGG AGGGTGGCTC TGTGAGGGGC ACTGGCTTCC TACAGGCCCC
4901 AGGTGGTCCT GAGGGGGCGG CTGCAAAGGC CAGGAGGCCC ACAGGCCCTT
4951 CTGCCCCTC CTGGGGAAGT GGATTTGGGG TCACTTTGTA TGAGGTGGGG

Figure 4 continued

6/21

5001 GCGGGTAACCA GCTTTGGGCC AAGCTGTCAC CCTGGATGGG CCATCACTTG
5051 CCTGCTCTGT ATAGGCCAGA TGGCCAGAAG CTGCTCCTGT CCTGTTGATG
5101 GCCCATCCTC GAGGTCTGGA CCCTCGGGAA GAGGAGCAGT TGGTGGCAGG
5151 GATGGGCCAC CGGAGACCTT CCTGACCTCC AGGACACGCA GCTGTGTGTG
5201 CCTGTCCCCA GGCCACATGC CACAGGGCTG GGGGCTCCTT GGGGCAGGGC
5251 TGGGCATTGG TCTGGCTACT CTTGGTATCG CCTCTGCCTC CCTGCCTCCC
5301 AGTCATCATC CTCCCACCTC TGCCTCCCTG CCTGTTCCCTC TCTTTCTCCT
5351 CAGGCCCTTC CGGACATTTC CTGCTCACCT AGGTCTGGGC AGGCGGGGTC
5401 AGGTGCCGGG TGTGAGCTCA CTCCTTCCGG CAGCAAGGTG TAGCTATGTG
5451 CCGGAAGGAA GGCCGCTGCT GTTGCTCCGC CTCTGAGTGC ATCCTTTCCA
5501 GGTCTCCAC ACTCCCCTGT GCCCCGACAC CTGGTGCCTC CTTCAGCCAT
5551 TGGTTCATGT GTCTCCAGG CACAGCTTTC TAGTCAGAG CCTCTAGGCT
5601 GGGTGCAGGA AGTGCTGAGG AAGTGGCAGC CGGGAGGCGA GCTGGCACCC
5651 TGTCCCTCCT GTTCTGTCC GTCCCTGGAG CTGGACCGTA TGGCCCCGCA
5701 TGTGTGATCC CCACCTGGGG CTGTGCCTCT GGGCAAGTTG GGAAGCTTGG
5751 TGAGCCTCAT TTTCATGTGC CCGCCTCCCA GTACTGATGT GCAGGTTGAA
5801 TGAGGTGCCA ACTGTAATGA GTTGAATGG CCCTGCTGGC TGGGTGGGAC
5851 TGGGGAGCAG GTGGGGGCCG CTGGGGGGCA CAGAGGCACA CCCAGTGCCT
5901 CAGTCAGGGA GAGGGTGACA GAGAAGCTCT GGGTGAGGCC CCACTCCAC
5951 TCTGGCCATG GCTGCTGCCC TTTGGTCCAC TGCAGTGAAC TGTGCCATGG
6001 GGCTGGACCT CTGTGGGGAT TGGTGGGCAG TGGGCTTTCT TCCCGCTTGG
6051 GGCTCTGAC CTCTGGGGGC AGGGCGCTGC CCGGTGGGA CAGTCGGAAG
6101 GCTGGTAGAG GGACCTGAGG GGTCTGTGTG GTGGCTGGGG GCAGGCCTCA
6151 GGAATTTGAC AGCAGGGATC TGGAAAAGCT TTAATAACAT TATTTGTTGT
6201 CAGGATTGGG AAATGCTCCC CTCCTCCCTC CCCCTCTTTC ATCTTAGAGA
6251 CTGCTGCACA TCTGGTCAGT GTGGTCTTCT TGGTGGCCCC CAAGGTGGCA
6301 GGGGTACAC TGTATGAAA CCGTCCCTG GGTATGTGGT GCAGACATGC
6351 ACATGCAGAT GGTGATTGGC AGGTTGTAGC ATGAGGTGGC TTTGGGACGG
6401 TTCCAGTGAC AGTGAGTGGG CTGGATCTGG GGGGTTCTGG GCAGTCCAT
6451 CAAGCGGATA CCCCCACAGA CTGTCCTCTT GGGATAGTTG GGCCTGGGAG
6501 CCCTGCTTGC CTTGCCAAAA GGCAGGCGCA GAGTCATGAA GAAGAGGGCT
6551 TGGGGGCTCA GAGCCCCACT GTGTGTGCAG CCCAGGGTGG ACCTGGAGGA
6601 GGTGCGTGGG CAGGCTGGGC CGGCCGGGGC CTGGGGTGGG GGGGCTGGT
6651 GTGGCAGGGA GGCAGGGCCA GACTGTCAGC GCTGCCTGGC TGAGGATGCT
6701 GGCACCCTGT CCTCCCCAGC CGTCTGTCTC CTGGGTGCAG CCATCTGAGT
6751 GCTGACCOCA GCCGCCCTG GAGGCTGGCT GTTCTCCTGT GCCCTATTGC
6801 TGGGGACATG TGCCACAGG AGGGAAAGG AAGGCCGGC CTCTCCGTT
6851 ~~ACAAAACCTG AGGCCTTCT CAATGCCCTG GATGGCCTCC TGGTGGCAGG~~
6901 GTGGTTGGTG GGAGGTGGGG CTGCTGCTTA GAACCCGCCA GCGGGCCTGG
6951 GCCTGGGCTG AGCTGCACCC CTCCAACCTC GCCTCCAGCT GAGGGTTGGC
7001 TTCCATCTCC ACCAGGCCCA GCACTGGGCA CAGGGCTCTC AGAGGCAGGC
7051 TCTGAAAGTC CCCTGCTGGC TTCTGCAGTG GACTCCAGGC GCCGAGCCCC
7101 CAGGGGGCTC GCATTGCGCT CACCCTGCGA AGCCACGTGA AGGCTGGGTC
7151 CTCCCCTCCG GAAGGGCCAA ATGCAGGGCA TGGGTGGTTT GAATGGTGGC
7201 CCCTGGGCTC CCCGGAGGGA CCAGCTGCTG TCGAGGGCCG CCCCCCTCCC
7251 ACTTCCGTCT TGCATACCA GCTCCTGTGG CACTCCCCAC GCCCGTCCC
7301 CCAGTGGGAG CGGCAGGCCC CCGGTGGCTC TGCCCGCGA GGGGATGTG
7351 TGGGCGGGG GGTGGCCTTG CTGCCAGATG CTCTGCCCCG AGTGTCCGTC
7401 TCCGCTCTCC AGTGTCCGTC CTCTGCCCCG AGTGTCCGTC
7451 TGGGCTGATG CTCTGCCCCG AGTGTCCGTC

Figure 4 continued

7/21

7501 GTACGTAGAG TGACACCTTG GAGCAAGGGT CCTGACGGCC GGGGGGCCAT
7551 GGGCTCTTCT CCAGGGGTAG GTGTCTGTAC TTGTGTAGCT GTGGTGAATG
7601 GAGCTCTGTG CTGGCGGTGG GGGTCCCTGG AGCAGCCGTA CCCTGGGACC
7651 CTACCGGGAG CATGCTCATG CCGTCCCTGC TGAATCCAG GAGATGCCTG
7701 CAGAGGGCAG CCTGGGAGCC TCTGAGCTGG GGTCTGCGCC CCAGGGGGCA
7751 CTGGAGTCTC CCCAGGGGGC GAGAGAGAGT AGGCAGGGAT GGTCTGGTGG
7801 CCCTGGGTGG GGGATGGCTG CTCCGTGGGC CCAGGCCCTC CCTGGCAGCA
7851 CAGGTGAGTG GTCTTGGGGG TCCACGTAGA ACTTCTCTT CTGTTCCAAA
7901 TTGCCCTCAT GGGTGCGGCA TGCCTGGGTG AACCTGGGGG AGCAGGGTGA
7951 GGACATGCTT CTCAGCCAG CCCACAGCTC CAGGCCACAC TCTGCAGGAC
8001 TCTGGCCOCT CCCTCAGCCC TGGAGGGAGC AGGACTGGAG TCCTGTGTCTC
8051 GCCTTGCTCT GACCTGGCCG AGGCCACTGC TGTGGGGCCC CAGCAGGCCT
8101 GCCCAGCAGA AGGTGGAGTG CAGGGACCCC AGGGGCAGCC TTCAGGGTGG
8151 GGCAGGGTGA GGCCCGACTG GGCCAGCCC CACCGCTCAG TGCTGATGTG
8201 GCGCGAGGCC TTCGCCOCTC CAGCTGACGT GTCTGCCTGC CCTGGGTGTG
8251 GCTCCAGAGG CTGCCTGTGT ACCAGGGGGC CCCAGCTTC TGTCTGTGGT
8301 TCTGGGCAGT CCCCTGGGGA GCGGTGGGGG CTGTGTGCCA GTCCAGACCC
8351 AGTAGTCCAC GCGTCTGGT CTCTGGAGGC CGTGGCTGGT CCAGGACTGT
8401 GGCAAGGTGG TCGTGCAGGG CAGCCOCTCA GCAGCTGTC TGTCTCTCTG
8451 CAGCCCCCAG CCTCCTGGCC CTTTGGTGCA CCCACAAAGC TCCOCCCTCC
8501 CCCAGGAGCT GGGGCCGCC TCTGCGTCTC CTCGGCAGCC TGGGCTTCCA
8551 GGTGGCTGGG CCTCTTTCGA GCTCCAACCTC TTGCCTGTGG TGGGCTCTCA
8601 GGACAGTCAA CTGCCAGTCG GCAGACATTG CAGGACCACG TGTGTCTTGG
8651 TAAGCTGGCT GGTTAGGTGT TTAGCTGGGG GATGGTGTGG CAGGTGGCCC
8701 CTGCATCTCT GAGCCTGTCA CCTCCTCGGG AAGCCTTCTG GGTGGGGGAC
8751 TCCACCCATG TCGCCTGGAG AAGCATCACT TTTCCACAGA GCCTTCTGCA
8801 ACCCCCGTGG GGCCTGAGCC TGGGGTGGGG GAGGTGGTGG CCCCTGCTCC
8851 TGCAGAGGCC AGCCAGGCAT CTGGCCOCCAG GCCACTGGCA AGAGCTCGTT
8901 GTGTTGGGGG ATCTGTCCTT TGCTGTGTGT GCAGGAGCGG CCGAGCCAGG
8951 CGGGGGCGTG AGTAGGGGTG GAGACCCAGG CCCAGCTTCC CCAGCCCTC
9001 AGGACCGGCC TGCTCTTTCC CACCACCCCA CCAAGTGCCT GGGCACACCC
9051 CGCCTGTGAG GATGGGCCCC GTTGGCAGGG CGGAGCCCTG GGAGGGTGGC
9101 AGTGCGCCGG GCAGGCTTGG ACTTCACTGG GGCTTGGGGT TGTCGTGTG
9151 GCCAGGGGCG CTGACCOGCT TGGTGGGACG GACGGCCGCT GGGCAGCAGG
9201 TTTCTTCTGC CACGGTGGCA CAGGCACCTG GGGTTGTGGT TGGCTCCAGG
9251 CGGGCGGGGG CTGCGTGCCC CTGCGCAGGC ACAAGGCGG TGGGTGGGGA
9301 GTCTCAGAGC TTGGCGTGAG GTCCCACAGG GCTGGCCCTG CAGGATGGAG
9351 GCCACTGTCC TGAGCTGCAG GTGCTGGCAG GAGCTGGGGT GGGCGTTCTG
9401 GGGCCGTGGC TGACAGOGTT ATGTCCCTCT CTCTCTATCG CAGTCTTAA
9451 ~~GCAGCTGAGC GGTCTTCTG GAGCTGAGT GGTGAGT GGGCCCCGGC~~
9501 CTGCCCCAGC CCCTGCCACC TCACCOCTCG CCTACACAGA CCCTCACCCA
9551 ~~CCTGCGTCTG CAGTCTGAGT GGTGAGT GGTGAGT GGTGAGT GGTGAGT~~
9601 ~~GTGAGTGGGT GGTGAGTGGT GGTGAGTGGT GGTGAGTGGT GGTGAGTGGT~~
9651 TGAGCTGGGT GCCCAGGAGG CCTCAGGCCG GCGGTGGGTG GGACAGGGCT
9701 GATCTGGGCC TGAACCTGCC CTGGGTGTCT TCTGTCCTCA GGTGAGT
9751 ~~GTGAGTGGGT GGTGAGTGGT GGTGAGTGGT GGTGAGTGGT GGTGAGTGGT~~ GTAAGC
9801 AGTGCCCTCA CGCCCTOCCC TGAATTGCCT CAAGGTCTTT ACCAGTCGGG
9851 CTTAGGGGGG GCCACCAGCT GGTCCCACTG TGCTTCAGGG TTTTGGGCCT
9901 TTCGTGGCCT TCCTGAGAGG GGCTGCACCT CAGGCCTGGT GGCTCTTCTC
9951 CAGGGAGGTC CTCTGACCAG GGAGGGGGGT CCCTGGCTGA CGCTCTGCTC

Figure 4 continued

8/21

10001 CCACCCCAG
10051
10101 GTGGG CCCCACCCCC GCCCCGCCCC CCGCCC CGC
10151 TGTCTCGGCC ACGGGCAGCG CGGGGGGCGT GGCCTGAGCT TGCTCTCCC
10201 ACAG
10251
10301 GT GAGGGCTGCC TCGGGCTGGG GCCACTGGGC
10351 TGCCACTTGC CTCGGGACCG GCAGGGGCTC GGCTCACCCC CGACCCGCCC
10401 CCTGCCGCTT GCTCGTAG
10451
10501 TCCTGO CTGGGGGG ACTGCCCGGC GGCTGGCCT GCTAGCCCCG
10551 CCCTCCCTTC CAG
10601
10651 GTG AGGCGGGGCC TCGTGGGCCA GGGTGGGCGG
10701 GCCTGCCGGC ACCCGGCACC GGGGCTCAGC TCACTGTCCG CTTGCTTCCT
10751 TCCCCAG
10801 GTGCCCGGGG GGGGGGGGGG GACTCTGGGG CCGTTGGGGA GCTGACTCTG
10851 CGCTTTTTC AG
10901 GTGAGC AGGCAGGCCT GGCAGGGTGG GTTCOGGGGT CAGGGCTGAG
10951 GGAGCCAGCT GTGCCCTGTG CCCACAG
11001 GTGAGTG CCTGCTGGGT GGGGACC GT
11051 GGGGGCGGGT GGGGCTGTTT TGGCACTGG CACCCACTCC CCACAG
11101
11151
11201 TGGGTGGCCT TGCCGGGGCG GGGGTGGTGG GGGCCCCCGC
11251 TGGGGCTGGG GCCGGAGCCC CTGCCACTC TGCCCCGCCC CCGCAG
11301
11351 GTGGGTGT GCGCCTGGGG GCGGGGGGTT GGGGGGTGGG
11401 ACGGGGTGC GTGGCCCGGC GCCCAGCCCA CTGCCGCTC CCCCAG
11451
11501 STCA GTGCACTGAG
11551 GGCGCGCCT GCCCCTGGTG GGGGTGGGGG TGGGGGTGGG GGCTCGCTGA
11601 CGCCCCTCTC CCCTCAG
11651 GTGAGCAGCC CTGGACCCCC
11701 GCTCCGCCCC GCCCGGAGAG CGCAGAGGCT CACTCCCGTC CTGTGTCCCC
11751 AG
11801
11851
11901
11951
12001
12051
12101
12151 AATAAA AGCTCCC TCAGCTCCA
12201 GGGCACAGGG CTGGCAGGAG GGGGCGGCC TCCCACGTGG GGCCATGCTG
12251 TGGGAAGGAG GCCCAGCGC CTGGAGAGGA GCTGGGGCTG TGGTGACCT
12301 CCCTGCCTCA CAGGGCTCTG TGGTCAGACG TCTTGCCCTG CAAGGTGGAG
12351 ACTCCATGCT CCAAGGCCCG CTGTGCTGA GGTCTGCACA CAAGTGGATT
12401 CAACTTGGGT CAGGCCAGAG GCTAAGGTGT GGAAGAGGGT TGAGAATCAG
12451 GCTGACTTGA ACGGCAGCAA AGACTCCAAG GCAAGGCTGC AGAGGTCTCA

Figure 4 continued

9/21

```

12251 TGGGAAGGAG GCCCCAGCGC CTGGAGAGGA GCTGGGGCTG TGGTGACCCCT
12301 CCCTGCCTCA CAGGGCTCTG TGGTCAGACG TCTTGCCCTG CAAGGTGGAG
12351 ACTCCATGCT CCAAGGCCCC CTGTGCCTGA GGTCTGCACA CAAGTGGATT
12401 CAACTTGGGT CAGGCCAGAG GCTAAGGTGT GGAAGAGGGT TGAGAATCAG
12451 GCTGACTTGA ACGGCAGCAA AGACTCCAAG GCAAGGCTGC AGAGGTCTCA
12501 GAGGCTATGC GCACAGTCCC CTGCTGGGGT GCTCACCTGG GCTGGGCTCT
12551 GGGCTGCTTG GACAAAGCAG GTGGCCTGGC TCAGCCCTCA CCGAGGGCCT
12601 CCCTTGGGGG CAGAGGTGGG CCTGATGCCA GGGGCTCCCC GTTTTTCAG
12651 GCCCTCAGCA GGTAGTTGGG TGTGGCCCTC AGGATACCTT GGTCCAGAG
12701 CTTGCCACTC AAAAAGCTTG GCAGTGAGGC AAGGGCAACC CCGGGCTGTT
12751 CCCCCCTCTA CTGGCTCTGC CGCCTGGGTT GGAAACCCTG AGGCTGTGCC
12801 AGGCAGGTGT ACCCTGACAG CCAGCCATGG CCCAGTAAGA TGGGTGCCCG
12851 AGGTGGTACC TGGGCAGCGG ACCCAGCTGT GCTGCCCCCG CCCAACCAG
12901 AAGCCGCTCT AGCCCATGGT GGTGCTCTGG GCGAGACAGG CTGGTTGGCT
12951 AGGCACTGTT TGGTCTACAG CAGGTGTAGG CAGCGTCTCC CTGACCCCTG
13001 CCTCCTAGGA AGCCACCACC CTGGGCCCTA CTCATCAGCA AGGACACGCA
13051 GCAGGGCTGA GCTGGGGGTG CGTGGGCTGC TACGGCCCGC CACCTCCATC
13101 ACATGCACCT CTGCACCCCC TGCTGCCTGA CTCAGGAGTG GGGGGGGGGG
13151 TCCTGTGCTT CCTTCACTCC AGACCCACGG TGCTGAQCCA GTGCACCCAC
13201 CTGGTCCTCT AGTGCGGACC TGGCCACAGG GCTCCTGTGG GCCACGCTG
13251 ATCCCGCCCT GGTCCCTTCA TAAAGAACTC TTAGACAT GCAGCCAGG
13301 GGAGCCAGGA GGCTCCAGTG TGCTGTGTCC ATCTGCCTCC CTCCAGCCCC
13351 TTCCGAGACA CTGCGCATCA TGCCCCCTC CACCCACC CACACTGGCA
13401 GGAGGAACAG ACAGGGAGAC CACACACAGA GCTCGTTGTT TATAAATCTC
13451 TGCCTGGCTC ATCGGTCTGT TTGTCCA
13501
13551 CACCTGCATC CTGGTCAGTC TGAGTGTGGC CGTGAAGCCC AGGTGAGCTG
13601 TGGCTCACAG GGCTAGGCCC TCGGTGCTGG CCGGGGGCCA CGCCCCACCC
13651 CCTCTCCCC CCTCCGCCAG CCAGGGGACC AGGCTCCTGG ACACCAGGCC
13701 TGCCCAAGGC CTGCTCTCCT CCTGGGGCTT
13751
13801
13851
13901
13951 GGGTCAAGGT GCCGGTGGG CCGGGGGGCG GGGTGGGGT GGGGGGCTCA
14001 GCTCAC
14051
14101 CAGTGAGGGC TCCCAGC

```

Highly identical to human sequence
SINE Sequence

Figure 4 continued

10/21




Figure 5

11/21




Figure 6

12/21




Figure 7

13/21




Figure 8

14/21

Ath	246	CIVWLKLVSYAHTSY.....DIRSIANAADK.....ANPE.....	↓	↓
Bna	227	CIVWLKLVSYAHTNY.....DIRTLANSSDK.....ANPE.....		
Pfr	252	CTVWLKLVSYAHTNY.....DIRVLAKSLOKWEAMSPYWNLD.....		
Cel	188	VIEALKEFISYGHVNYWARDARRKITELKTQVTDLAKKTCDEPKQFWDLDKDELSMH		
Mmu	211	SIMFLKLYSYRDVNLWCRR..V...KAKAVSTGKKVSG.....AAA		
Rno	211	SIIFLKLSYRDVNLWCRR..V...KAKAVSAGKKVSG.....AAA		
Cae	205	TILFLKLFYRDVNLWCRR.....A...RAKAASAGKRASS.....AAA		
Hsa	202	TILFLKLFYRDVNSWCRR.....A...RAKAASAGKKASS.....AAA		
Bta_1	199	TILFLKLFYRDVNLWCRRRAGA...KAKAALAGKKANG.....GAA		
Bta_2	199	TILFLKLFYRDVNLWCRRRAGA...KAKAALAGKAANG.....GAA		

Figure 9

15/21




Figure 10

16/21




Figure 11




Figure 12

18/21




Figure 13

19/21




Figure 14

20/21

Bull 899: 5-C-AA-A-C-C (HF)
6-G-GC-G-T-T

Bull 705: 6-C-AA-G-C-T (FV)
4-C-GC-G-T-T

↑ ↑ ↑

Figure 15

21/21




Figure 16

SEQUENCE LISTING

<110> Arbeitsgemeinschaft Deutscher Rinderzüchter e.V.

<120> Method of testing a mammal for its predisposition for
fat content of milk and/or its predisposition for meat
marbling

<130> F 1078 EP

<140>

<141>

<160> 4

<170> PatentIn Ver. 2.1

<210> 1

<211> 14117

<212> DNA

<213> Bos taurus

<400> 1

ctgccccgac	aggcctgaca	accaacaaca	agccttcctc	aatgccacta	gagaaatggg	60
aagtgcagac	cccttcctgc	agcctgcttt	ccacatcctg	acttccagat	tcagggggaca	120
tgtccccaca	ctgaggaggc	tttccttggt	agctggacca	ggctgggtgt	ggggaggaga	180
taccaagga	ataagaacct	cccatggcca	ccccagccc	ttaggctcta	gacaggggtga	240
gtcaagtga	gaagatgaat	ggcagggctg	tgctgggctc	agacaaccaa	ggaacataga	300
ctcctgcccc	agcaaagtgc	cttggttaacc	aggtaggtag	gcatgagcta	agaggctcca	360
aatctttgca	gacatgtggt	caaactggat	cagcccaggg	ccagcacagc	tgtctgcacc	420
ctggcagggg	acaggcccac	cagactccac	tgggtgtggac	agcaggaaag	cctgacctgc	480
agtagacctg	ctgcttcagg	gtgggatcac	ctgaggtggg	caccccttc	tggggagcac	540
tgtcagcctt	cataacctca	ggatgaaagc	ccccagtatt	ggtagagctt	aggtaggcat	600
cattgcccac	tctgcatatg	aagagtctga	ccctcagga	gagaagcagc	ttgccaaggg	660
ctgcctttga	cttaagccct	gctccagttg	ggcttccttg	gtggctcaga	ccctaaagaa	720
tctgcctgca	atgtgggaaa	cctgggttca	gtccctggga	cgggaagatc	ccctggagaa	780
gggatggcaa	cccactccag	tgttcttgcc	tgagaatccc	acggacagag	gagcctggcg	840
ggctgcagtc	catggagtcg	caaagagtcg	gacacgactg	agcaactaac	actttcactt	900
tctgcccac	tacccacccc	atctgaacct	gaatacctga	gtgggtccca	ctggcaggaa	960
gagaggctcc	tagaggccca	gtcctcccca	aggctcctca	gctttggggc	ctggattgac	1020
tgttccagga	ctctgatggg	cggctggggg	ggatgacggg	tagaggctgc	ctccccagtg	1080
actgggacag	gcctagcctt	gtctccacag	gtgtccatgg	acaggacttt	gcaatccaga	1140
ggatgggtgg	tgtggtgcag	gtgtgtgacc	actgtgtcca	gggtcttctc	tcacggggccc	1200
aaggcgcctc	caacctggag	tcagcccaag	gctctttcta	aatccccaaa	ccctccagc	1260
ccttcattcc	gccagcctgc	agattcctcg	tcccaagaca	gatgttgctt	ccaccagggg	1320
gagattcctc	attgagcttt	ctttcaacaa	ctcctcacgc	acatttgctc	ccaaaagacc	1380
ccacctatct	tgacgttttc	cctcgtgcct	cttcgctgtg	accctggcag	cacctcaatc	1440
aggatccaga	ggtaccaggg	ctgtaggccc	cgccctcccc	ggaggccccg	ccctccccgg	1500
aggccccgcc	ctccccggag	gccccgcctt	ccccggaggc	cccgcctcc	ccggaggccc	1560
cgcctgtat	caaccttgga	ccccgtcttc	ctcaaacagg	ccccgccccg	ccttggtaca	1620
gaggcccttc	ctgattggtg	ccttcacagt	ccgtgccttc	tcattggctt	gaggccctga	1680
tctctcaact	ccagcgggtg	aacccttggt	cctgcacgt	ccgggtcag	atcggtcttc	1740
tttgatgacc	ctcggcccac	cctgggtgtc	tcactccagc	tgtttcatgt	tagccgaagg	1800
caaaggagcc	tggacgcgga	cacaggggag	cgcacccaac	acgtaccttc	actcgtcagt	1860
ggctactgtg	ctcagcctct	ccaggccaac	aggcagcctg	agccgtcaat	cttctcctct	1920
gccaatcagc	gcgccagcca	ggctggccct	ctagtcaggg	ctcgggtactg	aaggatggca	1980
agtcccgaag	gtccccaggg	acgcgtgcgc	acgggttagg	gggcttccca	ccagctgcct	2040
gggagagggg	tagggagggg	aaggcagagc	tcccgggact	cagccctgct	gcgcgttcct	2100
gagaggactc	tctcctcctt	ccatcctccc	ttgggagcta	tactgagtc	tagcgtgag	2160
tggcccaact	ctgcctatga	atagacgaag	gtgcttgag	actggctaag	gggatactcc	2220
tgatccaccg	aggccggggc	tgtgaggagg	caagaggggt	tctccagcct	gatgaggtcg	2280

ctcgagccct	tccacacgct	actccaagac	acggggccagg	tagctccagc	ctgccaggta	2340
aggatgtcag	gctggcctca	gccgcaaagt	gtccagtgagg	agaacatgtc	accaggggtcc	2400
caggtgcctg	ttggttgagg	taagaggggtc	aggagcgagt	ccggcaggaa	ggaggcttga	2460
tctcaggctg	agcctcttgg	tttattttgct	ttcagagagg	cggctcttccc	agctttgtctt	2520
accccatggg	agtgaacgga	gtgggttctg	tggttagggg	tggttcttgt	gtaaaccagg	2580
cctaaactcc	cggngaaccc	tcgcattctg	agatccagga	tactcacact	ccatgctctt	2640
tgccaaatgt	ttgtgaaacc	aagtaagatc	ggccttgccc	gcgcacgggc	ctcactgtgc	2700
agttgttttg	gtgtattggt	tgcttcattc	aacgactgga	tgactgccga	ctgtgcaatg	2760
aaacagaaac	ctctgggtcc	ctgcgaatca	acaccccagg	atcctaactc	cctggcaaaa	2820
ctggcccaag	tggggaaggc	gggaagttct	gcaagtctgc	agatgaaggc	agaagcgggg	2880
cgggtggaga	ggcgggctgg	cttgtctact	gtgggggccc	gggcagggga	gaggtggcca	2940
ccctgggaat	aggtgggcat	ggcacaagtc	ccggaatgcg	aggactgcgg	cctttctccc	3000
cctccgttct	ctgacctggc	gcgtgtttga	acagcctaag	tggaggaaaa	gtgggtgcct	3060
acggtggtaa	ttagtgggtt	cacagagcac	gaccgtgccg	cgggatgtac	gttcggtaga	3120
cgcgttgggt	gtcagcctga	cgttaacgca	ctaggcattt	cataaataac	tacaacccca	3180
aattctgcgc	ctgagctgag	aaatgacgaa	atcctgtgtt	tatagagcgg	gacaaggggc	3240
aggcagcgg	cagcagaggc	ttgtttgcag	ctgcgccgaa	gccccgcgtg	ttcctcgtct	3300
gtccgggatt	gcatttgcca	ggagaccaca	actcccaggg	tgacccgcgc	gccagcggac	3360
tacaaaggta	tgccgcgccg	ggccctgggc	cagttagctg	ctccgggaac	tacgttccc	3420
aggactccga	gaggagccgt	ccggcacgga	tttgacgcg	ctgattggcg	gcgcggacca	3480
cggcagtggc	gtagtagagg	cgggtggcgg	agttggccaa	gggtccggag	gcggggccac	3540
aggcctcggg	tgctgccagc	ccggcgggct	acgacttggc	cgcggcgggg	tgcgaaactaa	3600
ggccatgggc	gaccgcggcg	gcgcggggcg	ctcccggcgc	cggaggacgg	ggtcgcggcc	3660
ttcgatccag	ggcggcagtg	ggcccgcggc	agcgggaagag	gaggtgcggg	atgtgggcgc	3720
cggaggggac	gcgcgggtcc	gggacacaga	caaggacgga	gacgtagacg	tgggcagcgc	3780
ccactgggac	ctgaggttagc	ggtgcgcgtg	acccttaacc	tttgacccct	gatacggggc	3840
ccctgcgacc	caacctgggtg	gcccaggcct	gtcggcggca	gctcgggctc	gagtcggaga	3900
gtctggcgcc	tggaaccttg	tgacacagctg	tgcccctcgg	gcctccacgg	ggaaacttag	3960
cgggaggttg	ggggcggagg	gtctcctgcc	cggaaacacc	aggtacgggg	gccgagggga	4020
gggcagcggc	tcaacttcta	gacgccctcc	ctctgccttc	ctttgggtggg	ttctgaagct	4080
ttcccagggt	gagcccaacta	cgcacagtg	cctctacctg	gaaggagata	caggggtcct	4140
tcttgagggc	tatgaggggt	gccttgtggg	ttgataaagc	tcccggggga	ggaggggtga	4200
ccggcggaga	acagaggcaga	gggcagtgcg	aggggatttc	tcacccctcg	cagacctcc	4260
agagaatgg	cttcacaaag	gtccctcatc	cgtcaccccg	cgattgactg	gcctaggatc	4320
ctgcttatta	ccagcacaaa	tggtctgctc	aggggtcaaa	tggttcctgt	aatgggaccc	4380
tcacccctgg	ttggggtaca	ggggaggagt	tggaagtgcg	cacaccacac	ggtgggcgcc	4440
ctgcttagct	gaaggactga	tggaaggag	ttgggggagc	aagctgcggc	tgaaagggag	4500
gatctgaccc	acgtgggcat	cagctaagtc	ctgctggctg	cctccaggcg	cccccttgc	4560
catectccac	gcccctcccc	ccagccctga	ccttcattct	ggtcaagggc	tctcaggggc	4620
tctggttttg	ggatcagctc	cagagctaga	ggttatcaag	gaggaagtgg	gcaacaggtc	4680
agtcagcaag	gatttgctat	cttactggg	tgctgtgggg	aggggaggga	caagggcagt	4740
tggggtgcag	gcactgtccc	tgcccttggg	gggcacacag	ttcacctgag	agataagata	4800
gcccgcagcc	tgaagagtga	gagcaaaagt	caggcacaga	gttcaggatg	acaccagggg	4860
aggggtggctc	tgtgaggggc	actggcttcc	tacaggcccc	aggtggctct	gagggggcgg	4920
ctgcaaaggc	caggaggccc	acaggccccct	ctgcccactc	ctgggggaact	ggatttgggg	4980
tcacttttga	tgaggtgggg	gcgggtacca	gctttggggc	aagctgtcac	cctggatggg	5040
ccatcacttg	cctgctctgt	ataggccaga	tggccagaag	ctgctcctgt	cctgttgatg	5100
gcccacctc	gaggtctgga	ccctcgggaa	gaggagcagt	tggtggcagg	gatggggccac	5160
cggagaccct	cctgacctcc	aggacacgca	gctgtgtgtg	cctgtcccca	ggccacatgc	5220
cacagggtcg	ggggcctcct	ggggcagggc	tgggcatttg	tctggctact	cttgggtatg	5280
cctctgcctg	cctgcctccc	agtcacatc	ctccacctc	tgccctcctg	cctgttctct	5340
tctttctcct	caggcccttc	cggacatttc	ctgctcacct	aggtctgggc	aggcgggggc	5400
aggtgcgggg	tgtgagctca	ctccttccgg	cagcaagggtg	tagctatgtg	ccggaaggaa	5460
ggcgcgtgct	gttgccctgc	ctctgagtgc	atcccttcca	ggtcctccac	actccctgt	5520
gccccgacac	ctgggtgcgtc	cttcagccat	tggttcatgt	gtcctccagg	cacagctttc	5580
tagtccagag	cctctaggct	gggtgcagga	agtgtctagg	aagtggcagc	cgggagggcg	5640
gctggcacc	tgctccctcct	tgctctgtcc	gtccctggag	ctggaccgta	tggccccgca	5700
tgtgtgatcc	ccacttgggg	ctgtgcctct	gggcaagttg	ggaagcttgg	tgagcctcat	5760
ttctcatgtg	ccgcctccca	gtactgatgt	gcagggtgaa	tgaggtgcca	actgtaatga	5820
gttggaaatg	ccctgctggc	tgggtgggac	tggggagcag	gtgggggccc	ctggggggca	5880
cagaggcaca	cccagtgcct	cagtcaggga	gaggggtgaca	gagaagctct	gggtgaggcc	5940

ccacctccac	tctggccatg	gctgctgccc	tttgggtccac	tgcagtgaac	tgtgccatgg	6000
ggctggacct	ctgtggggat	tgggtgggca	tgggctttct	tcccgtttgg	ggcctctgac	6060
ctctgggggc	agggcgctgc	ccgggtggga	cagtcggaag	gctggttagag	ggacctgagg	6120
ggtctgtgtg	gtggctgggg	gcaggcctca	ggaatttgac	agcagggatc	tggaaaagct	6180
ttaataacat	tatttggtgt	caggattggg	aaatgctccc	ctccccctc	cccctctttc	6240
atcttagaga	ctgctgcaca	tctggtcagt	gtggctcttc	tgggtggcccc	caaggtggca	6300
ggggtcacac	tgttatgaaa	ccgtcccctg	ggtagtggtt	gcagacatgc	acatgcagat	6360
ggtgattggc	aggttgtagc	atgaggtggc	tttgggacgg	ttccagtggc	agtgaagtgg	6420
ctggatctgg	gggggttctg	gcagggtccat	caagcggaata	ccccacaga	ctgtcctctt	6480
gggtagattg	ggcctgggag	ccctgcttgc	cttgccaaaa	ggcaggcgca	gagtcatgaa	6540
gaagagggct	tgggggctca	gagccccact	gtgtgtgcag	cccagggtgg	acctggagga	6600
ggtgcgtggg	caggctgggc	cggccggggc	ctgggggtgg	ggggcctggg	gtggcaggga	6660
ggcaggggca	gactgtcagc	gctgcctggc	tgaggatgct	ggcaccctgt	cctccccagc	6720
cgtctgtctc	ctgggtgcag	ccatctgagt	gctgacccca	gccgcccctg	gaggctggct	6780
gttctcctgt	gccctattgc	tggggacatg	tgtccacagg	agggaaaggg	aagccccggc	6840
ctctcccctt	acaaaactgg	aggccttgct	caatgccctg	gatggcctcc	tgggtggcagg	6900
gtgggttggg	ggaggtgggg	ctgctgctta	gaaccggcca	gcggggcctg	gcctgggctg	6960
agctgcaccc	ctccacctct	gcctccagct	gaggggttgg	ttccatctcc	accaggccca	7020
gcactgggca	cagggctctc	agaggcaggc	tctgaaagtc	ccctgctggc	ttctgcagtg	7080
gactccaggc	gccgagcccc	cagggggctc	gcattgcgct	caccctgcga	agccacgtga	7140
aggctgggtc	ctcccctccg	gaagggccaa	atgcagggca	tgggtgggtt	gaatgggtgg	7200
ccctgggctc	ccgggagggg	ccagctgctg	tgagggccgc	ccccctcccc	acttccgtct	7260
tgcatacaca	gtcctctgtg	cactccccac	gccccgtccc	ccagtgggag	cggcaggccc	7320
ccggtggctc	gtcccgcgga	gggggatgtg	tgggcggcgg	ggtggccttg	ctgccagatg	7380
ctctgccccg	agtgtccgtc	tccgctctcc	aggtgtcacc	gcctgcagga	ttccctgttc	7440
agtcttgaca	gtggcttcag	caactaccgt	ggcatcctga	attggtgtgt	ggtgatgtgt	7500
gtacgtagag	tgacaccttg	gagcaagggt	cctgacggcc	ggggggccat	gggctcttct	7560
ccaggggtag	gtgtctgtac	ttgtgtagct	gtgggtgaat	gagctctgtg	ctggcggttg	7620
gggtccctgg	agcagccgta	ccctgggacc	ctaccgggag	catgctcatg	ccgtccctgc	7680
tgaatcccag	gagatgcctg	cagagggcag	cctgggagcc	tctgagctgg	ggtctgcgcc	7740
ccagggggca	ctggagtctc	cccagggggc	gagagagagt	aggcagggat	ggtctggttg	7800
ccctgggtgg	gggatggctg	ctccgtgggc	ccaggccctc	cctggcagca	caggtgagtg	7860
gtcttggggg	tccacgtaga	acttctctct	ctgttccaaa	ttgccctcat	gggtgcggca	7920
tgcctgggtg	aacctggggg	agcagggtga	ggacatgctt	ctcagcccag	cccacagctc	7980
caggccacac	tctgcaggac	tctggccccct	ccctcagccc	tggaggggag	aggactggag	8040
tcctgtgtcc	gccttgctct	gacctggccg	aggccactgc	tgtggggccc	cagcaggcct	8100
gccagcaga	aggtggagtg	cagggacccc	aggggcagcc	ttcagggtgg	ggcagggtga	8160
ggcccagact	ggcccagccc	caccgctcag	tgctgatgtg	gcgcgaggcc	ttcgcccctc	8220
cagctgacgt	gtctgcctgc	cctgggtgtg	gctccagagg	ctgcctgtgt	accaggggcc	8280
cccacgcttc	tgtttgtggt	tctgggcagt	ccccggggga	gcggtggggg	ctgtgtgccca	8340
gtccagacct	agtagtccac	gcgtcctggt	ctctggaggc	cgtggctggt	ccaggactgt	8400
ggcaagttcc	tcgtgcaggg	caggccctca	gcagcctgtc	tgttctctct	cagccccagc	8460
cctcctggcc	ctttgggtga	cccacaaagc	tccccctccc	cccaggagct	ggggccgcct	8520
gctgcgtcct	ctcggcagcc	tgggcttcca	ggtggctggg	cctcttagca	gctccaactc	8580
ttgcctgtgg	tgggctctca	ggacaggcaa	ctgccagtcg	gcagacattg	caggaccacg	8640
tgtgtcctgg	taagctggct	ggttaggtgt	ttagctgggg	gatggtgtgg	caggtggccc	8700
ctgcatctct	gagcctgtca	cctcctcggg	aagccttctg	ggtgggggac	tccaccatg	8760
tcgcctggag	aagcatcact	tttccacaga	gccttctgca	acccccgtgg	ggcctgagcc	8820
tgggggtggg	gaggtgggtg	cccctgctcc	tgcagaggcc	agccaggcat	ctggccccag	8880
gccactggca	agagctcgtt	gtgttggggg	atctgtcctt	tgctgtctgt	gcaggagcgg	8940
ccgaggcagg	cgggggctg	agtaggggtg	gagaccagg	cccagcttcc	ccagcccctc	9000
aggaccggcc	tgctctttcc	caccacccca	ccaagtgcgt	gggcacaccc	gcctgtgtag	9060
gatggggccc	gttggcaggg	cggagccctg	ggaggggtgg	agtgcgcggg	gcaggcttgg	9120
acttcaactg	ggcttggggg	tgtcgctgtg	gccagggggc	ctgaccgcgt	tgggtgggac	9180
gacggccgct	gggcagcagg	tttcttctgc	cacggtggca	caggcacctg	gggttgtggt	9240
tggctccagg	cgggcggggg	ctgcgtgccc	ctgcgcaggc	acataggccg	tgggtgggga	9300
gtctcagagc	ttggcgtgag	gtcccacagg	gctgggcctg	caggatggag	gccactgtcc	9360
tgagctcctg	gtgctggcag	gagctggggg	gggcgttctg	gggcccgtgg	tgacagcgtt	9420
atgtccctct	ctctctatcg	cagatcttaa	gcaacgcacg	gttatttcta	gagaacctca	9480
tcaagtgaat	gggcccggcg	ctgccccacc	ccctgccacc	tcaccctcgc	cctacacaga	9540
ccctcaccca	cctgcgtctg	caggtatggc	atcctgggtg	accccatcca	ggtggtgtct	9600

ctgttcctga	aggacccta	cagctggcca	gctctgtgcc	tggtcattgg	tgagctgggt	9660
gcccaggagg	cctcaggccg	gcggtgggtg	ggacagggct	gatctgggcc	tgaacctgcc	9720
ctgggttgct	tctgtcctca	gtggccaata	tctttgccgt	ggctgcgttc	caggtggaga	9780
agcgcctggc	cgtggtaagc	agtgcctca	cgccctcccc	tgacttgctt	caaggtcctt	9840
accagtcggg	cttagggcgg	gccaccagct	ggtcccactg	tgtctcaggg	ttttgggcct	9900
ttcgtggcct	tcctgagagg	ggctgcacct	caggcctggt	ggctcttctt	caggggaggtc	9960
ctctgaccag	ggaggggggt	ccctggctga	cgctctgtct	ccaccccagg	gagctctgac	10020
ggagcaggcg	gggctgtctg	tgcacggggt	caacctggcc	accattctct	gcttcccagc	10080
ggcctgtggc	tttctcctcg	agtctatcac	tccagggtgg	ccccaccccc	gccccgcgcc	10140
ccgcccacgc	tgtctcggcc	acgggcagcg	cggggggctg	ggcctgagct	tgcctctccc	10200
acagtgggct	ccgtgctggc	cctgatgggt	tacaccatcc	tcttctctca	gctgttctcc	10260
taccgggacg	tcaacctctg	gtgccgagag	cgcagggtct	gggccaaggc	caaggctggt	10320
gagggctgcc	tcgggctggg	gccactgggc	tgccacttgc	ctcgggaccg	gcaggggctc	10380
ggctcacccc	cgaccgcgcc	cctgccgctt	gctcgtagct	ttggcaggta	aggcggccaa	10440
cgggggagct	gcccagcgca	ccgtgagcta	ccccgacaac	ctgacctacc	gcggtgagga	10500
tcttgccggg	ggctgggggg	actgcccggc	ggcctggcct	gctagccccc	ccctcccttc	10560
cagatctcta	ctacttctct	ttcgccccca	ccctgtgcta	cgagctcaac	ttcccccgct	10620
ccccccgcat	ccgaaagcgc	ttcctgctgc	ggcgactcct	ggagatgggt	aggcggggcc	10680
tcgtgggcca	gggtgggcgg	gcctgccggc	acccggcacc	ggggctcagc	tcactgtccg	10740
cttgcttctt	tccccagctg	ttcctcaccc	agctccaggt	ggggctgata	cagcagggtac	10800
gtgcccgggg	gggggggggg	gactctgggg	ccgttggggg	gctgactctg	cgctttttgc	10860
agtggatggg	ccgggccatc	cagaactcca	tgaagccctt	caagggtgagc	aggcaggcct	10920
ggcagggtgg	gttcgggggt	cagggctgag	ggagccagct	gtgcccgtgt	cccacaggac	10980
atggactact	ccgcctcgt	ggagcgcctc	ctgaagctgg	cggtgagtgg	ctgctggggt	11040
ggggacgcgt	gggggcgggt	ggggctgttc	tggcacctgg	cacccactcc	ccacagggtc	11100
ccaaccacct	catctggctc	atcttcttct	actggctctt	ccactcctgc	ctgaacgcgc	11160
tggctgagct	catgcagttt	ggagaccgcg	agtctctacc	ggactgggtg	tgggtggcct	11220
tgcgggggcg	gggggtggtg	gggccccgcg	tggggctggg	gccggagccc	ctgcccactc	11280
tgccccgcgc	ccgcaggaac	tccgagtcca	tcacctactt	ctggcagaa	tggaaacatcc	11340
ctgttcacaa	gtggtgcata	aggtgggtgt	gcgcctgggg	gcgggggggt	gggggggtgg	11400
acggggctgc	gtggcccggc	gcccagccca	ctgcgcctc	ccccgcagac	acttctacaa	11460
gcccctgctc	cggcggggca	gcagcaagtg	ggcagccagg	acggcagtg	ttctggcctc	11520
cgccctcttc	cacgaggta	gtgcactgag	ggcgcgcctt	gcccctggtg	gggggtgggg	11580
tgggggtggg	ggctcgctga	cgccccctct	ccctcagtac	ctgggtgagca	tccccctggg	11640
aatgttccgc	ctctgggcct	tcaccggcat	gatggcgag	gtgagcagcc	ctggaccccc	11700
gctccgcccc	gccccgcgag	cgcagaggct	cactcccgtc	ctgtgtcccc	agatccccgt	11760
ggcctggata	gtggggccgt	tcttccgcgg	caactacggc	aacgcggccg	tgtggctgtc	11820
actcatcatc	gggcagccgg	tggccgtcct	gatgtacgtc	cacgactact	acgtgtctaa	11880
ccgtgagggg	ccggcagccg	gcacctgagc	gcctccagge	tggccccctc	gtgggtgttg	11940
gactgctttg	cgcgctgcc	tgcggctgga	ctagagcctg	ccccaacctg	ggtgcagcag	12000
gaggaggcct	ggctgggtga	agctgcctcc	tggcctccac	caggcctctg	ctcgaaggcg	12060
ttctctctgc	caggggagag	caggccccgac	gcagttcttg	ccccctgggag	gtgcccctgc	12120
tctggaaacc	ctacagatct	cgcccaaggg	tctgaatgtg	tcaataaagt	gctgtgcaca	12180
gtgagctccc	tcagcctcca	gggcacaggg	ctggcaggag	ggggcgggcc	tcccacgtgg	12240
ggccatgctg	tgggaaggag	gccccagcgc	ctggagagga	gctggggctg	tgggtgacct	12300
ccctgcctca	cagggtctct	tggctcagacg	tcttgccctg	caagggtggag	actccatgct	12360
ccaaggcccc	ctgtgcctga	ggtctgcaca	caagtggatt	caacttgggt	caggccagag	12420
gctaagggtg	ggaagagggt	tgagaatcag	gctgacttga	acggcagcaa	agactccaag	12480
ccaaggctgc	agaggctctca	gaggctatgc	gcacagtccc	ctgctggggg	gctcacctgg	12540
gctgggctct	gggctgcttg	gacaaagcag	gtggcctggc	tcagccctca	ccgagggcct	12600
cccttggggg	cagagggttg	cctgatgccca	ggggctcccc	gtttttccag	gcccctcagca	12660
ggtagtgggg	tgtggccctc	aggatacctt	ggtcccagag	cttgccactc	aaaaagcttg	12720
gcagtgaggg	aaggggcaacc	ccgggctggt	ccccctctca	ctggctctgc	cgccctgggt	12780
ggaaaccctg	aggctgtgcc	aggcagggtgt	accctgacag	ccagccatgg	cccagtaaga	12840
tgggtgcccc	agggtgtacc	tgggcagcgg	accagctgt	gctgcccccg	ccccaccag	12900
aagccgctct	agcccaggtg	ggtcgtctgg	gcgagacagg	ctgggtgggt	aggcactgtt	12960
tgggtctacg	caggtgtagg	cagcgtctcc	ctgacccctg	cctcctagga	agccaccacc	13020
ctgggccccta	ctcatcagca	aggacagcga	gcagggtgta	gctgggggtg	ctggtggctgc	13080
tacggccccg	cacctccatc	acatgcacct	ctgcaccccc	tgtgtgctga	ctcaggagtg	13140
gggggggggg	tctgtgtctt	ccttctactcc	agaccacagg	tgtgtaccca	gtgcacccac	13200
ctggtcctct	agtgcggacc	tggccacagg	gctcctgtgg	gcccacgctg	atccccccct	13260


```

ggtccttca taaagaactc ttgagcacat gcagcccagg ggagccagga ggctccagtg 13320
tgctgtgtcc atctgcctcc ctccagcccc ttccgagaca ctgcgcatca tgccccctc 13380
caccaccacc cacactggca ggaggaacag acagggagac cacacacaga gctcgttgtt 13440
tataaatctc tgcttggtc atcggtctgt ttgtccatgt atatatctgt atatctctat 13500
ggaaggggaa agggggactc gtgtaaaaat ccaaaatata attctatgaa cacctgcac 13560
ctggtcagtc tgagtgtggc cgtgaagccc aggtgagctg tggctcacag ggctaggccc 13620
tcggtgctgg ccggggggcca cgccccaccc cctctcccc cctccgccag ccaggggacc 13680
aggctcctgg acaccaggcc tgcccaaggc ctgctctcct cctggggctt ctacgagaca 13740
gtgggggtcct tggctttggg gggttctgag cccgtcagca gggagatggg ggggtcatcc 13800
gagtagtcgt ctccctcgga gaagtaggag ccctcccca gctcgaagag caccggcagg 13860
tcgctgctcc ccacgtccac ggagcccggg tccaggagca gcaggggctg ggcggtgtag 13920
tgcaccaact gcttccttag ggggtgcgact gggtaaggt gccggtgggg ccggggggcg 13980
gggtgggggt ggggggctca gctcacctga gtctgggctg cttttctctg cctccagagg 14040
tctggggggc tctgggggag agaggagctc ctggatctgc tggggcagca ggagggagca 14100
cagtgagggc tcccgcg

```

<210> 2

<211> 489

<212> PRT

<213> Bos taurus

<400> 2

```

Met Gly Asp Arg Gly Gly Ala Gly Gly Ser Arg Arg Arg Arg Thr Gly
  1                      5                      10                      15

Ser Arg Pro Ser Ile Gln Gly Gly Ser Gly Pro Ala Ala Ala Glu Glu
          20                      25                      30

Glu Val Arg Asp Val Gly Ala Gly Gly Asp Ala Pro Val Arg Asp Thr
          35                      40                      45

Asp Lys Asp Gly Asp Val Asp Val Gly Ser Gly His Trp Asn Leu Arg
          50                      55                      60

Cys His Arg Leu Gln Asp Ser Leu Phe Ser Ser Asp Ser Gly Phe Ser
          65                      70                      75                      80

Asn Tyr Arg Gly Ile Leu Asn Trp Cys Val Val Met Leu Ile Leu Ser
          85                      90                      95

Asn Ala Arg Leu Phe Leu Glu Asn Leu Ile Lys Tyr Gly Ile Leu Val
          100                      105                      110

Asp Pro Ile Gln Val Val Ser Leu Phe Leu Lys Asp Pro Tyr Ser Trp
          115                      120                      125

Pro Ala Leu Cys Leu Val Ile Val Ala Asn Ile Phe Ala Val Ala Ala
          130                      135                      140

Phe Gln Val Glu Lys Arg Leu Ala Val Gly Ala Leu Thr Glu Gln Ala
          145                      150                      155                      160

Gly Leu Leu Leu His Gly Val Asn Leu Ala Thr Ile Leu Cys Phe Pro
          165                      170                      175

Ala Ala Val Ala Phe Leu Leu Glu Ser Ile Thr Pro Val Gly Ser Val
          180                      185                      190

Leu Ala Leu Met Val Tyr Thr Ile Leu Phe Leu Lys Leu Phe Ser Tyr
          195                      200                      205

```

Arg Asp Val Asn Leu Trp Cys Arg Glu Arg Arg Ala Gly Ala Lys Ala
 210 215 220
 Lys Ala Ala Leu Ala Gly Lys Ala Ala Asn Gly Gly Ala Ala Gln Arg
 225 230 235 240
 Thr Val Ser Tyr Pro Asp Asn Leu Thr Tyr Arg Asp Leu Tyr Tyr Phe
 245 250 255
 Leu Phe Ala Pro Thr Leu Cys Tyr Glu Leu Asn Phe Pro Arg Ser Pro
 260 265 270
 Arg Ile Arg Lys Arg Phe Leu Leu Arg Arg Leu Leu Glu Met Leu Phe
 275 280 285
 Leu Thr Gln Leu Gln Val Gly Leu Ile Gln Gln Trp Met Val Pro Ala
 290 295 300
 Ile Gln Asn Ser Met Lys Pro Phe Lys Asp Met Asp Tyr Ser Arg Ile
 305 310 315 320
 Val Glu Arg Leu Leu Lys Leu Ala Val Pro Asn His Leu Ile Trp Leu
 325 330 335
 Ile Phe Phe Tyr Trp Leu Phe His Ser Cys Leu Asn Ala Val Ala Glu
 340 345 350
 Leu Met Gln Phe Gly Asp Arg Glu Phe Tyr Arg Asp Trp Trp Asn Ser
 355 360 365
 Glu Ser Ile Thr Tyr Phe Trp Gln Asn Trp Asn Ile Pro Val His Lys
 370 375 380
 Trp Gly Ile Arg His Phe Tyr Lys Pro Met Leu Arg Arg Gly Ser Ser
 385 390 395 400
 Lys Trp Ala Ala Arg Thr Ala Val Phe Leu Ala Ser Ala Phe Phe His
 405 410 415
 Glu Tyr Leu Val Ser Ile Pro Leu Arg Met Phe Arg Leu Trp Ala Phe
 420 425 430
 Thr Gly Met Met Ala Gln Ile Pro Leu Ala Trp Ile Val Gly Arg Phe
 435 440 445
 Phe Arg Gly Asn Tyr Gly Asn Ala Ala Val Trp Leu Ser Leu Ile Ile
 450 455 460
 Gly Gln Pro Val Ala Val Leu Met Tyr Val His Asp Tyr Tyr Val Leu
 465 470 475 480
 Asn Arg Glu Ala Pro Ala Ala Gly Thr
 485

<210> 3

<211> 14117

<212> DNA

<213> Bos taurus

<400> 3

ctgccccgac	aggcctgaca	accaacaaca	agccttcctc	aatgccacta	gagaaatggg	60
aagtgcagac	cccttcctgc	agcctgcttt	ccacatcctg	acttccagat	tcaggggaca	120
tgtccccaca	ctgaggaggc	tttccttggt	agctggacca	ggctggttgt	ggggaggaga	180
taccaagga	ataagaacct	cccatggcca	ccccagccc	ttaggctcta	gacagggatga	240
gtcaagttga	gaagatgaat	ggcagggctg	tgctgggctc	agacaacca	ggaacataga	300
ctcctgcccc	agcaaagcc	cttggttaacc	aggtaggtag	gcatgagcta	agaggctcca	360
aatcttttga	gacatgtggt	caaactggat	cagcccaggg	ccagcacagc	tgtctgcacc	420
ctggcagggg	acaggcccac	cagactccac	tggtgtggac	agcaggaaag	cctgacctgc	480
agtagacctg	ctgcttcagg	gtgggatcac	ctgaggtggg	caccccttc	tggggagcac	540
tgtagcctt	cataacctca	ggatgaaagc	ccccagatt	ggtagagctt	aggtaggcat	600
cattgcccc	tctgcatatg	aagagtctga	ccctcagggg	gagaagcagc	ttgccaaggg	660
ctgcctttga	cttaagccct	gctccagttg	ggcttccttg	gtggctcaga	ccctaaagaa	720
tctgcctgca	atgtgggaaa	cctgggttca	gtccctggga	cggaagatc	ccctggagaa	780
gggatggcaa	ccactccag	tgttcttgcc	tgagaatccc	acggacagag	gagcctggcg	840
ggctgcagtc	catggagtcg	caaagagtcg	gacacgactg	agcaactaac	actttcactt	900
tctgccccaa	tacccacccc	atctgaacct	gaatacctga	gtgggtccca	ctggcaggaa	960
gagaggctcc	tagaggccca	gtcctcccca	aggctcctca	gctttggggc	ctggattgac	1020
tgttccagga	ctctgatggg	cggctggggg	ggatgacggg	tagaggctgc	ctccccagtg	1080
actgggacag	gcctagcctt	gtctccacag	gtgtccatgg	acaggacttt	gcaatccaga	1140
ggatgggtgg	tggtgtgcag	gctgctgacc	actgtgtcca	gggtcttctc	tcacggggcc	1200
aaggcgcctc	caacctggag	tcagcccaag	gctctttcta	aatccccaaa	cccttcacgc	1260
ccttcattcc	ccagcctgc	agattcctcg	tcccaagaca	gatgttgctt	ccaccagggg	1320
gagattcctc	attgagcttt	ctttcaacaa	ctcctcacgc	acatttgctc	ccaaaagacc	1380
ccacctatct	tgacgttttc	cctcgtgcct	cttcgctgtg	accctggcag	cacctcaatc	1440
aggatccaga	ggtaccaggg	ctgtaggccc	cgccctcccc	ggaggccccg	ccctccccgg	1500
aggccccgcc	ctccccggag	gccccgccct	ccccggaggc	cccgccctcc	ccggaggccc	1560
cgccctgtat	caaccttgga	ccccgtcttc	ctcaaacagg	ccccgccccg	ccttggtaca	1620
gaggcccttc	ctgattgggtg	ccttcacagt	ccgtgccttc	tcattggctt	gaggccctga	1680
tctctcaact	ccagcgggtg	aacctttggt	tccttcacgt	ccgggtcag	atcggttctc	1740
tttgatgacc	ctgggccac	cctggtgtcc	tcactccagc	tgtttcatgt	tagccgaagg	1800
caaaggagcc	tggaacggga	cacagggagc	cgcccccaac	acgtaccttc	actcgtcagt	1860
ggctactgtg	ctcagcctct	ccaggccaac	agccagcctg	agccgtcaat	ctctcctctt	1920
gccaatcagc	gcgccagcca	ggctggccct	ctagtcaggg	ctcggtactg	aaggatggca	1980
agtcccgaag	gctcccaggg	acgcgtgcgc	acgggttagg	gggcttccca	ccagctgcct	2040
gggagaggga	tagggaggga	aaggcagagc	tcccgggact	cagccctgct	gcgcgttctt	2100
gagaggactc	tctcctcctt	ccatcctccc	ttgggagcta	tactgagtc	tagcgtgag	2160
tggcccaact	ctgcctatga	atagacgaag	gtgcttggac	actggctaag	gggatactcc	2220
tgatccaccg	aggccgggccc	tgtgaggagg	caagaggggt	tctccagcct	gatgaggctg	2280
ctcgagccct	tccacacgct	actccaagac	acgggcccagg	tagctccagc	ctgccaggta	2340
aggatgtcag	gctggcctca	gcccgaatg	gtccagtggg	agaacatgtc	accaggggtcc	2400
caggtgcctg	ttggttgagg	taagaggggtc	aggagcgagt	ccggcaggaa	ggaggcttga	2460
tctcaggctg	agcctcttgg	tttatttgct	ttcagagagg	cggtcttccc	agcttctgctt	2520
accccatggg	agtgaacgga	gtgggttctg	tggttagggg	tgtttcttgt	gtaaaccagg	2580
cctaaactcc	cgggtgaacc	tcgatctctg	agatccagga	tactcacact	ccatgctctt	2640
tgccaaatgt	ttgtgaaacc	aagtaagatc	ggccttgccc	gcgcacgggc	ctcactgtgc	2700
agttgtttt	gtgtattggt	tgcttcattc	aacgactgga	tgactgccga	ctgtgcaatg	2760
aaacagaaac	ctctgggtcc	ctgcgaatca	acaccccagg	atcctaactc	cctggcaaaa	2820
ctggcccaag	tggggaaggc	gggaagttct	gcaagctgc	agatgaaggc	agaagcgggg	2880
cggttgagg	ggcgggctgg	cctgtctact	gtgggggccc	gggcagggga	gaggtggcca	2940
ccctgggaat	aggtgggcat	ggcacaagtc	ccggaatgcg	aggactgcgg	ccttctctcc	3000
cctccgttct	ctgacctggc	gcgtgtttga	acagcctaag	tggaggaaaa	gtgggtgcct	3060
acggttggt	ttagtgggtt	cacagagcac	gaccgtgccg	cggtatgtac	gttcggtaga	3120
cgcgttggt	gtcagcctga	cgttaacgca	ctaggtcatt	cataaataac	tacaacccca	3180
aattctgcgc	ctgagctgag	aatgacgaa	atcctgtgtt	tatagagcgg	gacaaggggc	3240
agccagcgg	cagcagaggc	ttgtttgcag	ctgcccggaa	gccccgcgtg	ttcctcgtct	3300
gtccgggatt	gcatttgcca	ggagaccaca	actcccagg	tgcaccgcgc	gccagcggac	3360
tacaaaggta	tgccgcgcgc	ggccctgggc	cagttagctg	ctccgggaac	tacgttctcc	3420
aggactccga	gaggagccgt	ccggcacgga	tttgacgcg	ctgattggcg	gcgggacca	3480
cgccagtgcc	gtagtagagg	cgggtggcgg	agttggccaa	gggtccggag	gcggggccac	3540
aggcctcggg	tgctgccagc	ccggcgggct	acgacttggc	cgcgccgggg	tgccaactaa	3600

ggccatgggc	gaccgcggcg	gcgcggggcg	ctcccggcgc	cggaggacgg	ggtcgcggcc	3660
ttcgatccag	ggcggcagtg	ggcccgcggc	agcggaaagag	gaggtgcggg	atgtggggcg	3720
cggaggggac	gcgcgggtcc	gggacacaga	caaggacgga	gacgtagacg	tgggcagcgg	3780
ccactgggac	ctgaggtagc	ggtgcgcgtg	acccctaacc	tttgaccctc	gatacggggc	3840
ccctgcgacc	caacctgggtg	gcccaggcct	gtcggcgcca	gctcgggctc	gagtcggaga	3900
gtctggcgcc	tggaccttgg	tgcacagctg	tgcccctcgg	gcctccacgg	ggaaacttag	3960
cgggaggttg	ggggcggagg	gtctcctgcc	cggaaacacc	aggtacgggg	gccgagggga	4020
gggcagcggc	tcaacttcta	gacgccttcc	ctctgccttc	ctttggtggg	ttctgaagct	4080
ttcccagggt	gagccacta	cgcacagtgt	cctctacctg	gaaggagata	caggggtcct	4140
tcctgagggc	tatgaggggt	gccttgtggg	ttgataaagc	tcccggggga	ggaggggtga	4200
cggcgggaga	acagaggcag	gggcagtgcg	aggggatttc	tcatccctcg	cagaccctcc	4260
agagaatggt	cttcacaaa	gtccctcatc	cgtcacccgg	cgattgactg	gcctaggatc	4320
ctgcttatta	ccagcacaaa	tggctgctct	agggctcaa	tgggtcctgt	aatgggaccc	4380
tcaccctgg	ttgggggtaca	ggggaggagt	tggaaagtgc	cacaccaca	ggtgggcggc	4440
ctgcttagct	gaaggactga	tgggaaggag	ttgggggagc	aagctgcggc	tgaaagggag	4500
gatctgacct	acgtgggcat	cagctaagtc	ctgctggctg	cctccaggcg	ccccctttgc	4560
catcctccac	gccccctccc	ccagccctga	ccttcctcct	ggtcaagggc	tctcaggggc	4620
tctgggtttg	ggatcagctc	cagagctaga	ggttatcaag	gaggaagtgg	gcaacaggtc	4680
agtcagcaag	gatttgctat	cttcactggg	tgctgtgggg	aggggagggg	caagggcagt	4740
tgggggtgcag	gcactgtccc	tgcccttggg	gggcacacag	ttcacctgag	agataagata	4800
gccgcagccc	tgaagagtga	gagcaaaggt	caggcacaga	gttcaggatg	acaccagggg	4860
aggggtggctc	tgtgaggggc	actggcttcc	tacaggcccc	aggtgggtcct	gagggggcgg	4920
ctgcaaaggg	caggagggcc	acaggccctc	ctgccactc	ctggggaact	ggattttggg	4980
tcactttgta	tgaggtgggg	gcgggtacca	gctttgggcc	aagctgtcac	cctggatggg	5040
ccatcaactg	cctgctctgt	ataggccaga	tggccagaag	ctgctcctgt	cctgttgatg	5100
gcccatectc	gaggtctgga	ccctcgggaa	gaggagcagt	tgggtggcagg	gatggggccac	5160
cggagaccct	cctgacctcc	aggacacgca	gctgtgtgtg	cctgtcccca	ggccacatgc	5220
cacagggctg	ggggcctcct	ggggcagggc	tgggcattgg	tctggctact	cttggtatcg	5280
cctctgcctc	cctgcctccc	agtcatcatc	ctcccacctc	tgccctccctg	cctgttcctc	5340
tctttctcct	caggcccttc	cggacatttc	ctgctcacct	aggtctgggg	aggcggggtc	5400
aggtgcgggg	tgtgagctca	ctccttcggg	cagcaagggtg	tagctatgtg	ccggaaggaa	5460
ggccgctgct	gttgccctgc	ctctgagtgc	atcccttcca	ggtccctccac	actcccctgt	5520
gccccgacac	ctgggtgcgtc	cttcagccat	tggttcatgt	gtccctccagg	cacagctttc	5580
tagtccagag	cctctaggct	gggtgcagga	agtgtctgagg	aagtggcagc	cgggagggcg	5640
gctggcaccc	tgtccctcct	tgttctgtcc	gtccctggag	ctggaccgta	tggccccgca	5700
tgtgtgatcc	ccacttgggg	ctgtgcctct	gggcaggttg	ggaagcttgg	tgagcctcat	5760
tttcatgtgc	ccgcctccca	gtactgatgt	gcaggttgaa	tgaggtgcca	actgtaatga	5820
gttggaaatgg	ccctgctggc	tgggtgggac	tggggagcag	gtggggggccg	ctgggggggca	5880
cagaggcaca	cccagtgcct	cagtcaggga	gaggggtgaca	gagaagctct	gggtgaggcc	5940
ccacctccac	tctggccatg	gctgctgccc	tttgggtccac	tgcaagtgaac	tgtgccatgg	6000
ggctggacct	ctgtggggat	tgggtggcag	tgggctttct	tcccgttgg	ggcctctgac	6060
ctctgggggc	agggcgctgc	ccgggtggga	cagtcgggaag	gctggtagag	ggacctgagg	6120
ggtctgtgtg	gtggctgggg	gcaggcctca	ggaatttgac	agcagggatc	tggaaaagct	6180
ttaataacat	tatttgttgt	caggattggg	aaatgtctcc	ctccccctc	ccccctcttc	6240
atcttagaga	ctgctgcaca	tctgggtcagt	gtggtcttct	tgggtggccc	caaggtggca	6300
ggggtcacac	tgttatgaaa	ccgtccccctg	ggtatgtggg	gcagacatgc	acatgcagat	6360
ggtgattggc	aggttgtagc	atgaggtggc	tttgggacgg	ttccagtga	agtgagtggt	6420
ctggatcttg	ggggttcttg	gcaggctccat	caagcggata	ccccacaga	ctgtcctctt	6480
gggatatgtg	ggcctgggag	ccctgcttgc	cttgccaaaa	ggcaggcgca	gagtcatgaa	6540
gaagagggct	tgggggctca	gagccccact	gtgtgtgcag	cccaggggtg	acctggagga	6600
ggtgcgtggg	caggctgggc	cggccggggc	ctgggggtgg	ggggcctggg	gtggcagggg	6660
ggcagggcca	gactgtcagc	gctgcctggc	tgaggatgct	ggcaccctgt	cctccccagc	6720
cgtctgtctc	ctgggtgcag	ccatctgagt	gctgaccca	gcccggcctg	gaggtgggt	6780
gttctcctgt	gccctattgc	tggggacatg	tgtccacagg	agggaaagg	aagccccggc	6840
ctctccccct	acaaaactgg	aggccttgct	caatgccctg	gatggcctcc	tgggtggcagg	6900
gtggttggtg	ggaggtgggg	ctgctgctta	gaaccgcgca	gcgggcctgg	gcctgggctg	6960
agctgcaccc	ctccacctct	gcctccagct	gaggggtggc	ttccatctcc	accaggccca	7020
gcactgggca	cagggtctct	agaggcagct	cttgaaagtc	ccctgctggc	ttctgcagt	7080
gactccaggc	gccgagcccc	cagggggctc	gcattgcgct	caccctgcga	agccacgtga	7140
aggctgggtc	ctccccctcg	gaagggccaa	atgcagggca	tgggtgggtt	gaatgggtgg	7200
ccctgggctc	cccggaggga	ccagctgctg	tgagggccgc	ccccctcccc	acttccgtct	7260

tgcataacca	gtcctctgtg	cactccccac	gccccgtccc	ccagtgggag	cggcaggccc	7320
ccgggtggctc	tgcccccgga	gggggatgtg	tgggcggcgg	ggtggccttg	ctgccagatg	7380
ctctgccccg	agtgtccgtc	tccgctctcc	aggtgtcacc	gcctgcagga	ttccctgttc	7440
agttctgaca	gtggcttcag	caactaccgt	ggcatcctga	attgggtgtg	ggtgatgtctg	7500
gtacgtagag	tgacaccttg	gagcaagggt	cctgacggcc	ggggggccat	gggctcttct	7560
ccaggggtag	gtgtctgtac	ttgtgtagct	gtgggtgaatg	gagctctgtg	ctggcgggtg	7620
gggtcccttg	agcagccgta	ccctgggacc	ctaccgggag	catgctcatg	ccgtccctgc	7680
tgaatcccag	gagatgcctg	cagagggcag	cctgggagcc	tctgagctgg	ggtctgcgcc	7740
ccagggggca	ctggagtctc	cccagggggc	gagagagagt	aggcagggat	ggtctgggtg	7800
ccctgggttg	gggatggctg	ctccgtgggc	ccaggccctc	cctggcagca	caggtagtg	7860
gtcttggggg	tccacgtaga	acttccctct	ctgttccaaa	ttgccctcat	gggtgcggca	7920
tgccctgggtg	aacctggggg	agcaggggtg	ggacatgctt	ctcagcccag	cccacagctc	7980
caggccacac	tctgcaggac	tctggcccc	ccctcagccc	tggagggagc	aggactggag	8040
tcctgtgtcc	gccttgctct	gacctggccg	aggccactgc	tgtggggccc	cagcaggcct	8100
gcccagcaga	aggtggagtg	cagggacccc	aggggcagcc	ttcagggtgg	ggcagggtga	8160
ggcccagactg	ggcccagccc	caccgctcag	tgctgatgtg	gcgcgaggcc	ttcggccctc	8220
cagctgacgt	gtctgcctgc	cctgggtgtg	gctccagagg	ctgcctgtgt	accagggggc	8280
cccacgcttc	tgtttgtggt	tctgggcagt	cccctgggga	gcggtggggg	ctgtgtgcca	8340
gtccagaccc	agtagtccac	gcgtcctggt	ctctggaggc	cgtggctggt	ccaggactgt	8400
ggcaaggttg	tcgtgcaggg	caggccctca	gcagccctgc	tgttctcctg	cagccccag	8460
cctcctggcc	ctttgggtga	cccacaaagc	tccccctccc	cccaggagct	ggggccgcct	8520
gtgcgtccct	ctcgccagcc	tgggcttcca	ggtggctggg	cctcttagca	gctccaactc	8580
ttgcctgtgg	tgggctctca	ggacaggcaa	ctgccagtcg	gcagacattg	caggaccacg	8640
tgtgtcctgg	taagctggct	ggttaggttg	ttagctgggg	gatgggtgtg	caggtagccc	8700
ctgcctctct	gagcctgtca	cctcctcggg	aagcctctctg	ggtgggggac	tccaccctatg	8760
tcgcctggag	aagcatcact	tttccacaga	gccttctgca	acccccgtgg	ggcctgagcc	8820
tgggggtggg	gaggtgggtg	cccctgctcc	tgcagaggcc	agccaggcat	ctggccccag	8880
gccactggca	agagctcggt	gtgttggggg	atctgtcctt	tgctgtctgt	gcaggagcgg	8940
ccgaggcagg	cgggggctgt	agtaggggtg	gagacccagg	cccagcttcc	ccagccctc	9000
aggaccggcc	tgctctttcc	caccacccca	ccaagtgcgt	gggcacaccc	cgcctgtgag	9060
gatggcccg	ggtggcaggg	cggagccctg	ggagggtggc	agtgcgcggg	gcaggcttgg	9120
acttctactg	ggcttggggg	tgtcgtctgt	gccaggggcg	ctgaccgctg	tgggtgggacg	9180
gacggccgct	gggcagcagg	tttcttctgc	cacggtggca	caggcacctg	gggttgtggt	9240
tggctccagg	cgggcggggg	ctgcgtgccc	ctgcgcaggc	acataggccg	tgggtgggga	9300
gtctcagagc	ttggcgtgag	gtcccacagg	gctgggcctg	caggatggag	gccactgtcc	9360
tgagctgcag	gtgctggcag	gagctggggg	gggcgttctg	gggccgtggc	tgacagcggt	9420
atgtccctct	ctctctatcg	cagatcttaa	gcaacgcacg	gttatttcta	gagaacctca	9480
tcaagtgagt	gggccccggc	ctgccccagc	ccctgccacc	tcaccctcctg	cctacacaga	9540
ccctcaccca	cctgcgtctg	caggatggc	atcctggtgg	accccatcca	ggtggtgtct	9600
ctgttcttga	aggacccta	cagctggcca	gctctgtgcc	tggtcattgg	tgagctgggt	9660
gcccaggagg	cctcaggccg	gcggtgggtg	ggacagggct	gatctgggac	tgaacctggc	9720
ctgggttgct	tctgtcctca	gtggccaata	tctttgccgt	ggctgcgttc	caggtaggga	9780
agcgctggc	cgtggtaagc	agtgcctca	cgcctccccc	tgacttgctt	caaggctcct	9840
accagtcggg	cttagggcgg	gccaccagct	ggtccactg	tgtctcaggg	ttttgggcct	9900
ttcgtggcct	tcttgagagg	ggctgcacct	caggcctggt	ggctcttctt	cagggagggtc	9960
ctctgaccag	ggaggggggt	ccctggctga	cgctctgctc	ccaccccagg	gagctctgac	10020
ggagcaggcg	gggctgctgc	tgcacggggg	caacctggcc	accattctct	gcttcccagc	10080
ggccgtggcc	tttctcctcg	agtctatcac	tccagggtgg	ccccaccccc	gccccgcgcc	10140
ccgcccacgc	tgtctcgccc	acgggcagcg	cggggggcgt	ggcctgagct	tgccctctcc	10200
acagtgggct	ccgtgctggc	cctgatggtc	tacaccatcc	tcttctcctca	gctgttctct	10260
taccgggacg	tcaacctctg	gtgccgagag	cgcagggctg	gggccaaggc	caaggctgggt	10320
gagggctgcc	tcgggctggg	gccactgggc	tgccacttgc	ctcgggaccg	gcaggggctc	10380
ggctcaccac	cgaccgcgcc	cctgccgctt	gctcgtagct	ttggcaggta	agaaggccaa	10440
cgggggagct	gcccagcgca	ccgtgagcta	ccccgacaac	ctgacctacc	gcggtgagga	10500
tcctgccggg	ggctgggggg	actgcccggc	ggcctggcct	gctagccccg	ccctcccttc	10560
cagatctcta	ctaactctct	ttcgccccca	ccctgtgcta	cgagctcaac	ttcccccgct	10620
ccccccgcat	ccgaaagcgc	ttcctgctgc	ggcgactcct	ggagatgggt	aggcggggcc	10680
tcgtggggcca	gggtggggcg	gcctgcggcc	acccggcacc	ggggctcagc	tcactgtccg	10740
cttgccttct	tccccagctg	ttcctcacc	agctccaggt	ggggctgac	cagcaggtac	10800
gtgcccgggg	gggggggggg	gactctgggg	ccgttggggg	gctgactctg	cgttttttgc	10860
agtggatggt	cccggccatc	cagaactcca	tgaagccctt	caaggtagag	aggcaggcct	10920

```

ggcaggggtgg gttccgggggt cagggctgag ggagccagct gtgccctgtg cccacaggac 10980
atggactact cccgcacatcgt ggagcgcctc ctgaagctgg cggtagtggt cctgctgggt 11040
ggggacgcgt gggggcgggt ggggctgttc tggcacctgg caccactcc ccacagggtcc 11100
ccaaccacct catctggctc atcttcttct actggctctt ccactcctgc ctgaacgccg 11160
tggctgagct catgcagttt ggagaccgag agttctaccg ggactgggtg tgggtggcct 11220
tgccggggcg ggggtgggtg gggcccccgc tggggctggg gccggagccc ctgcccactc 11280
tgccccgccc ccgcaggaac tccgagtcca tcacctactt ctggcagaac tggaaacatcc 11340
ctgttcacaa gtggtgcac cagggtgggtg gcgcctgggg gcgggggggt ggggggtggg 11400
acgggggtgc gtggcccggc gccagccca ctgccgcctc ccccgagac acttctacaa 11460
gcccattgctc cggcggggca gcagcaagtg ggcagccagg acggcagtggt ttctggcctc 11520
cgccttcttc cacgaggtca gtgcactgag ggcgcgcctt gccctgggtg ggggtggggg 11580
tgggggtggg ggctcgtgta cgcctctctc ccctcagtag ctgggtgagca tccccctggg 11640
aatgttccgc ctctgggcct tcaccggcat gatggcgag gtgagcagcc ctggaccccc 11700
gtcccgcccc gcccgcgag cgcagaggct cactccgctc ctgtgtcccc agatcccgct 11760
ggcctggata gtgggcccgt tcttccgagg caactacggc aacgcggccg tgtggctgtc 11820
actcatcatc gggcagccgg tggcgtcct gatgtacgtc cactcctcag cactcctcag 11880
ccgtgaggcg ccggcagccg gcacctgagc gcctccaggc tggccccctc gtgggtgttg 11940
gactgctttg ccgcgctgcc tgcggctgga ctgagcctg cccaacctg ggtgcagcag 12000
gaggaggcct ggctgggtgga agctgcctcc tggcctccac caggcctctg cctgaagggt 12060
ttctcctctc caggggagag cagggccgac gcagttcttg cccctgggag gtgcccattg 12120
tctggaaacc ctacagatct cgcaccaagg tctgaatgtg tcaataaagt gctgtgcaca 12180
gtgagctccc tcagcctcca gggcacaggg ctggcaggag ggggcggccc tcccactgtg 12240
ggccatgctg tgggaaggag gccccagcgc ctgggagga gctggggctg tggtgacctg 12300
ccctgcctca cagggtctct tggtcagacg tcttgccctg caagggtggg actccatgct 12360
ccaaggcccc ctgtgcctga ggtctgcaca caagtggatt caacttgggt caggccagag 12420
gctaagggtg ggaagagggt tgagaatcag gctgacttga acggcagcaa agactccaag 12480
gcaaggctgc agaggtctca gaggctatgc gcacagtccc ctgctggggg gctcacctgg 12540
gctgggctct gggctgcttg gacaaagcag gtggcctggc tcagccctca ccgagggcct 12600
cccttggggg cagaggttg cctgatgcca ggggctcccc gtttttccag gccctcagca 12660
ggtagttggg tgtggccctc aggatacctt ggtcccagag ctggccactc aaaaagcttg 12720
gcagtgaggc aagggaacc cgggctgtt cccccctcta ctggctctgc cgcctgggtt 12780
ggaaaccctg aggcgtgtgc aggcaggtgt accctgacag ccagccatgg cccagtaaga 12840
tgggtgcccc aggtgggtacc tgggcagcgg acccagctgt gctgcccccg cccaaccag 12900
aagccgctct agcccatggg ggtcgtctgg gcgagacagg ctggttgggt aggcactgtt 12960
tgggtctacag caggtgtagg cagcgtctcc ctgaccctg cctcctagga agccaccacc 13020
ctgggcccta ctcacagca aggacagcga gcagggctga gctgggggtg cgtgggctgc 13080
tacggcccg caccctccatc acatgcacct ctgcaccccc tgctgctga ctcaggagtg 13140
gggggggggg tctgtgtgtt ccttcaactc agacccacgg tgctgacca gtgcacccac 13200
ctggtcctct agtgcgacc tggccacagg gctcctgtgg gccacgctg atcccgccct 13260
ggtcctctta taaagaactc ttgagcacat gcagcccagg ggagccagga ggtccagtg 13320
tgctgtgtcc atctgcctcc ctccagcccc ttccgagaca ctgcgcatca tgccccctc 13380
cacccccacc cacactggca ggaggaacag acagggagac cacacacaga gctcgttgtt 13440
tataaatctc tgcctggctc atcggctgtt ttgtccatgt atatatctgt atatctctat 13500
ggaaggggaa agggggactc gtgtaaaaat ccaaaatata attctatgaa cacctgcac 13560
ctggtcagtc tgagtgtggc cgtgaagccc aggtgagctg tggctcacag ggctaggccc 13620
tcggtgctgg ccgggggcca cgcgccaccc cctctcccc cctccgccag ccaggggacc 13680
aggctcctgg acaccaggcc tgcccaaggc ctgctctcct cctggggctt ctacgagaca 13740
gtgggtcctt tggcttggg ggggtctgag cccgtcagca gggagatggg ggggtcatcc 13800
gagtagtcgt ctcctcgga gaagtaggag cctcccccga gctcgaagag caccggcagg 13860
tcgctgctcc ccacgtccac ggagcccggg tccaggagca gcaggggctg ggcggtgtag 13920
tgcaccaact gcttccctag ggggtgcgact ggggtcaagg ggcggtgggg ccggggggcg 13980
gggtgggggt ggggggctca gctcacctga gtctgggctg cttttctctg cctccagagg 14040
tctggggggc tcctggggag agaggagctc ctggatctgc tggggcagca ggagggagca 14100
cagtgagggc tcccgcg 14117

```

<210> 4

<211> 489

<212> PRT

<213> Bos taurus

<400> 4
Met Gly Asp Arg Gly Gly Ala Gly Gly Ser Arg Arg Arg Arg Thr Gly
1 5 10 15
Ser Arg Pro Ser Ile Gln Gly Gly Ser Gly Pro Ala Ala Ala Glu Glu
20 25 30
Glu Val Arg Asp Val Gly Ala Gly Gly Asp Ala Pro Val Arg Asp Thr
35 40 45
Asp Lys Asp Gly Asp Val Asp Val Gly Ser Gly His Trp Asn Leu Arg
50 55 60
Cys His Arg Leu Gln Asp Ser Leu Phe Ser Ser Asp Ser Gly Phe Ser
65 70 75 80
Asn Tyr Arg Gly Ile Leu Asn Trp Cys Val Val Met Leu Ile Leu Ser
85 90 95
Asn Ala Arg Leu Phe Leu Glu Asn Leu Ile Lys Tyr Gly Ile Leu Val
100 105 110
Asp Pro Ile Gln Val Val Ser Leu Phe Leu Lys Asp Pro Tyr Ser Trp
115 120 125
Pro Ala Leu Cys Leu Val Ile Val Ala Asn Ile Phe Ala Val Ala Ala
130 135 140
Phe Gln Val Glu Lys Arg Leu Ala Val Gly Ala Leu Thr Glu Gln Ala
145 150 155 160
Gly Leu Leu Leu His Gly Val Asn Leu Ala Thr Ile Leu Cys Phe Pro
165 170 175
Ala Ala Val Ala Phe Leu Leu Glu Ser Ile Thr Pro Val Gly Ser Val
180 185 190
Leu Ala Leu Met Val Tyr Thr Ile Leu Phe Leu Lys Leu Phe Ser Tyr
195 200 205
Arg Asp Val Asn Leu Trp Cys Arg Glu Arg Arg Ala Gly Ala Lys Ala
210 215 220
Lys Ala Ala Leu Ala Gly Lys Lys Ala Asn Gly Gly Ala Ala Gln Arg
225 230 235 240
Thr Val Ser Tyr Pro Asp Asn Leu Thr Tyr Arg Asp Leu Tyr Tyr Phe
245 250 255
Leu Phe Ala Pro Thr Leu Cys Tyr Glu Leu Asn Phe Pro Arg Ser Pro
260 265 270
Arg Ile Arg Lys Arg Phe Leu Leu Arg Arg Leu Leu Glu Met Leu Phe
275 280 285
Leu Thr Gln Leu Gln Val Gly Leu Ile Gln Gln Trp Met Val Pro Ala
290 295 300
Ile Gln Asn Ser Met Lys Pro Phe Lys Asp Met Asp Tyr Ser Arg Ile
305 310 315 320

SUBSTITUTE SHEET (RULE 26)

12

Val Glu Arg Leu Leu Lys Leu Ala Val Pro Asn His Leu Ile Trp Leu
 325 330 335
 Ile Phe Phe Tyr Trp Leu Phe His Ser Cys Leu Asn Ala Val Ala Glu
 340 345 350
 Leu Met Gln Phe Gly Asp Arg Glu Phe Tyr Arg Asp Trp Trp Asn Ser
 355 360 365
 Glu Ser Ile Thr Tyr Phe Trp Gln Asn Trp Asn Ile Pro Val His Lys
 370 375 380
 Trp Gly Ile Arg His Phe Tyr Lys Pro Met Leu Arg Arg Gly Ser Ser
 385 390 395 400
 Lys Trp Ala Ala Arg Thr Ala Val Phe Leu Ala Ser Ala Phe Phe His
 405 410 415
 Glu Tyr Leu Val Ser Ile Pro Leu Arg Met Phe Arg Leu Trp Ala Phe
 420 425 430
 Thr Gly Met Met Ala Gln Ile Pro Leu Ala Trp Ile Val Gly Arg Phe
 435 440 445
 Phe Arg Gly Asn Tyr Gly Asn Ala Ala Val Trp Leu Ser Leu Ile Ile
 450 455 460
 Gly Gln Pro Val Ala Val Leu Met Tyr Val His Asp Tyr Tyr Val Leu
 465 470 475 480
 Asn Arg Glu Ala Pro Ala Ala Gly Thr
 485

SUBSTITUTE SHEET (RULE 26)